Index of Authors

ÅBERG, L. *see* MALMSTRÖM, A. 477–489

ABRAHAM, E. P. *see* FAWCETT, P. A. 741–746; USHER, J. J. 729–739

ALLEN, G. & HARRIS, J. I. The binding of nicotinamide–adenine dinucleotide to glyceraldehyde 3-phosphate dehydrogenase from *Bacillus stearothermophilus* 747–749

AMBLER, R. P. The amino acid sequence of *Staphylococcus aureus* penicillinase 197–218

AMPHLETT, G. W. *see* JACKSON, P. 85–97

ANDERSON, A. J. & ARCHIBAND, A. R. Poly(glucosylglycerol phosphate) teichoic acid in the walls of *Bacillus stearothermophilus* B65 115–120

ARCHIBALD, A. R. *see* ANDERSON, A. J. 115–120

AZHAR, S. & MENON, K. M. J. Adenosine 3′:5′-cyclic monophosphate-dependent and plasma-membrane-associated protein kinase(s) from bovine corpus luteum. Properties of associated enzyme and phosphorylation of specific plasma-membrane proteins 23–36

BACCINO, F. M., ZURETTI, M. F. & PERNIGOTTI, L. Stimulation of rat liver β-galactosidase activity by ions 567–573

BADDILEY, J. *see* LAMBERT, P. A. 671–676; POWELL, D. A. 387–397

BARCLAY, A. N., LETARTE-MUIRHEAD, M. & WILLIAMS, A. F. Purification of the Thy-1 molecule from rat brain 699–706

BARCLAY, A. N. *see also* LETARTE-MUIRHEAD, M. 685–697

BARRETT, M. C. & DAWSON, A. P. The reaction of choline dehydrogenase with some electron acceptors 677–683

BETTS, S. A. & MAYER, R. J. Purification and properties of 6-phosphogluconate dehydrogenase from rabbit mammary gland 263–270

BLAIR, A. H. *see* SIDHU, R. S. 443–445

BOHUON, C. *see* BOUDENE, C. 413–415

BOLAND, M. J. & GUTFREUND, H. Pig heart lactate dehydrogenase. Binding of pyruvate and the interconversion of pyruvate-containing ternary complexes 715–727

BOUDENE, C., DUPREY, F. & BOHUON, C. Radioimmunoassay of colchicine 413–415

BRENNA, O., PERRELLA, M., PACE, M. & PIETTA, P. G. Affinity-chromatography purification of alkaline phosphatase from calf intestine 291–296

BRINKWORTH, R. I., MASTERS, C. J. & WINZOR, D. J. Evaluation of equilibrium constants for the interaction of lactate dehydrogenase isoenzymes with reduced nicotinamide–adenine dinucleotide by affinity chromatography 631–636

BROCKLEHURST, K. *see* STUCHBURY, T. 417–432

BROWN, D. G. *see* BROWN, K. B. 505–512

BROWN, K. B., NELSON, N. F. & BROWN, D. G. Effects of polyamines and methylglyoxal bis(guanylhydrazone) on hepatic nuclear structure and deoxyribonucleic acid template activity 505–512

BRUNORI, M., PARR, S. R., GREENWOOD, C. & WILSON, M. T. A temperature-jump study of the reaction between azurin and cytochrome *c* oxidase from *Pseudomonas aeruginosa* 185–188

CAMMACK, R. *see* MULLINGER, R. N. 75–83

CARLSTEDT, I. *see* MALMSTRÖM, A. 477–489

CHEN, S.-S. & ENGEL, P. C. Dogfish M_4 lactate dehydrogenase: reversible inactivation by pyridoxal 5′-phosphate and complete protection in complexes that mimic the active ternary complex 447–449

CHEN, S.-S. & ENGEL, P. C. Reversible modification of pig heart mitochondrial malate dehydrogenase by pyridoxal 5′-phosphate 297–303

CHEN, S.-S. *see also* ENGEL, P. C. 305–318

CHERAYIL, J. D. *see* THIMMAPPAYA, B. 377–386

CHOI, H. U. & MEYER, K. The structure of keratan sulphates from various sources 543–553

CITTANOVA, N. *see* LAURENT, C. 513–518

CLAMP, J. R. *see* KAMERLING, J. P. 491–495

COLBY, J., DALTON, H. & WHITTENBURY, R. An improved assay for bacterial methane mono-oxygenase. Some properties of the enzyme from *Methylomonas methanica* 459–462

CORNISH-BOWDEN, A. *see* MARGISON, G. P. 249–256; STORER, A. C. 361–367

CULLIS, P. R. *see* SIM, E. 555–560

DALTON, H. *see* COLBY, J. 459–462

DARLISON, M. G. *see* STORER, A. C. 361–367

DAWSON, A. P. *see* BARRETT, M. C. 677–683

DEAN, R. T. *see* MESSER, M. 17–22

DE LAUZON, S. *see* LAURENT, C. 513–518

DE LEY, M. & OSAKI, S. Intramolecular electron transport in human ferroxidase (caeruloplasmin) 561–566

DESAI, R. *see* HARWOOD, J. L. 707–714

DEVERSON, E. V. *see* MILSTEIN, C. P. 615–624

DICKSON, D. P. E. *see* MULLINGER, R. N. 75–83

DIETRICH, C. P. *see* MICHELACCI, Y. M. 121–129

DIXON, H. B. F. Factorization of the Michaelis functions 271–274

DON, M. M., MASTERS, C. J. & WINZOR, D. J. Further evidence for the concept of bovine plasma arylesterase as a lipoprotein 625–630

DOWBEN, R. M. *see* WARD, P. E. 755–758

DOYEN, N. *see* LAPRESLE, C. 637–643

DUCKWORTH, M. *see* POWELL, D. A. 387–397

DUPREY, F. *see* BOUDENE, C. 413–415

EDWARDS, R. G., THOMAS, P. & WESTWOOD, J. H. The purification and properties of a β-*N*-acetylhexosaminidase from *Trichomonas foetus* 145–148

ENGEL, P. C. & CHEN, S.-S. A product-inhibition study of bovine liver glutamate dehydrogenase 305–318

ENGEL, P. C. *see also* CHEN, S.-S. 297–303, 447–449

ENGELBORGHS, Y., MARSH, A. & GUTFREUND, H. A quenched-flow study of the reaction catalysed by creatine kinase 47–50

ENNS, C. A. *see* KANESHIRO, C. M. 433–434

ERDÖS, E. G. *see* WARD, P. E. 755–758

EYRE, D. R. & MUIR, H. The distribution of different molecular species of collagen in fibrous, elastic and hyaline cartilages of the pig 595–602

FAWCETT, P. A., USHER, J. J. & ABRAHAM, E. P. Behaviour of tritium-labelled isopenicillin N and 6-aminopenicillanic acid as potential penicillin precursors in an extract of *penicillium chrysogenum* 741–746
FEINSTEIN, A. *see* MILSTEIN, C. P. 615–624
FRANSSON, L.-Å. *see* MALMSTRÖM, A. 477–489
FREARSON, N. & PERRY, S. V. Phosphorylation of the light-chain components of myosin from cardiac and red skeletal muscles 99–107
FRIEDEN, E. *see* OSAKI, S. 519–525

GAUCHER, G. M. *see* STEVENSON, K. J. 527–542
GEDNEY, C. D., *see* WARD, P. E. 755–758
GEISOW, M. J. An improved method for purifying sialidase 181–183
GERWIG, G. J. *see* KAMERLING, J. P. 491–495
GOLL, D. E. *see* HAMMOND, K. S. 189–192
GOODGAME, D. M. L., JEEVES, I., REYNOLDS, C. D. & SKAPSKI, A. C. Heavy metal–pyrimidine nucleotide interaction. X-ray structure of a cadmium derivative of cytidine 5′-monophosphate 467–468
GRAZI, E. Fructose 1,6-diphosphate aldolase from rabbit muscle. Effect of pH on the rate of formation and on the equilibrium concentration of the carbanion intermediate 167–172
GREEN, M. L. Crystallization of one of the chicken pepsinogens and the derived pepsin 763–764
GREEN, M. L. *see also* LLEWELLIN, J. M. 319–326
GREENWOOD, C. *see* BRUNORI, M. 185–188; PARR, S. R. 51–59
GUTFREUND, H. *see* BOLAND, M. J. 715–727; ENGELBORGHS, Y. 47–50

HAHN, M. G. *see* KANESHIRO, C. M. 433–434
HALBAN, P. A. & OFFORD, R. E. The preparation of a semisynthetic tritiated insulin with a specific radioactivity of up to 20 curies per millimole 219–225
HALL, D. O. *see* MULLINGER, R. N. 75–83
HAMMOND, K. S. & GOLL, D. E. Purification of insect myosin and α-actinin 189–192
HANCOCK, I. C. *see* LAMBERT, P. A. 671–676
HARFORD, S. & WEITZMAN, P. D. J. Evidence for isosteric and allosteric nucleotide inhibition of citrate synthase from multiple-inhibition studies 455–458
HARRIS, J. I. *see* ALLEN, G. 747–749
HARWOOD, J. L., DESAI, R., HEXT, P., TETLEY, T. & RICHARDS, R. Characterization of pulmonary surfactant from ox, rabbit, rat and sheep 707–714
HAY, A. W. M. The transport of 25-hydroxycholecalciferol in a new world monkey 193–196
HEINEGÅRD, D. *see* THYBERG, J. 157–166
HERBERT, J. A. L. *see* STUCHBURY, T. 417–432
HEXT, P. *see* HARWOOD, J. L. 707–714
HICKMAN, P. E. & WEIDEMANN, M. J. The purification and properties of pig spleen phosphofructokinase 327–336
HOLLAWAY, M. R. *see* OSBORNE, H. H. 37–45
HOOGHWINKEL, G. J. M. *see* OVERDIJK, B. 257–261
HOPWOOD, J. J. & ROBINSON, H. C. Studies on the polydispersity and heterogeneity of cartilage proteolgycans. Identification of three proteoglycan structures in bovine nasal cartilage 581–594

IDLER, W. W. *see* STEINERT, P. M. 603–614
IUPAC COMMISSION ON THE NOMENCLATURE OF ORGANIC CHEMISTRY & IUPAC–IUB COMMISSION ON BIOCHEMICAL NOMENCLATURE. Nomenclature of carotenoids. (Recommendations 1974) 5–7
IUPAC–IUB COMMISSION ON BIOCHEMICAL NOMENCLATURE. The nomenclature of peptide hormones. Recommendations (1974) 1–4

JACKSON, P., AMPHLETT, G. W. & PERRY, S. V. The primary structure of troponin T and the interaction with tropomyosin 85–97
JÄNNE, J. *see* PÖSÖ, H. 67–73
JAYLE, M.-F. *see* LAURENT, C. 513–518
JEEVES, I. *see* GOODGAME, D. M. L. 467–468
JELLINCK, P. H. *see* MCNABB, T. 275–279
JOHNSON, C. E. *see* MULLINGER, R. N. 75–83
JOHNSON, T. C. *see* TAUB, F. 173–180
JONES, M. N. A theoretical approach to the binding of amphipathic molecules to globular proteins 109–114

KAMERLING, J. P., GERWIG, G. J., VLIEGENTHART, J. F. G. & CLAMP, J. R. Characterization by gas–liquid chromatography–mass spectrometry and proton-magnetic-resonance spectroscopy of pertrimethylsilyl methyl glycosides obtained in the methanolysis of glycoproteins and glycopeptides 491–495
KANESHIRO, C. M., ENNS, C. A., HAHN, M. D., PETERSON, J. S. & REITHEL, F. J. Evidence for an active dimer of *Escherichia coli* β-galactosidase 433–434
KEIR, H. M. *see* MCLENNAN, A. G. 227–238, 239–247
KELLIE, A. E. *see* SAMARAJEEWA, P. 369–376
KIDO, R. *see* NOGUCHI, T. 399–406
KITSON, T. M. The effect of disulfiram on the aldehyde dehydrogenases of sheep liver 407–412
KRALL, J. F., SOCHER, S. H., VAN, N. T. & O'MALLEY, B. W. Fractionation of chick oviduct chromatin. Nuclease-resistant deoxyribonucleic acid 497–503

LAMBERT, P. A., HANCOCK, I. C. & BADDILEY, J. Influence of alanyl ester residues on the binding of magnesium ions to teichoic acids 671–676
LAPRESLE, C. & DOYEN, N. Isolation and properties of a fragment of human serum albumin demonstrating the absence of a methionine residue from some of the albumin molecules 637–643
LARA, F. J. S. & OKRETIC, M. C. A micro-method for the assay of polyadenylate-containing ribonucleic acid by gel electrophoresis 575–580
LAURENT, C., DE LAUZON, S., CITTANOVA, N., NUNEZ, E. & JAYLE, M.-F. The comparative specificity of three oestradiol-binding proteins. Rat α-foetoprotein, rat liver 17β-hydroxy steroid dehydrogenase and anti-(oestradiol-6-carboxymethyloxime–bovine serum albumin) antiserum 513–518
LETARTE-MUIRHEAD, M., BARCLAY, A. N. & WILLIAMS, A. F. Purification of the Thy-1 molecule, a major cell-surface glycoprotein of rat thymocytes 685–697
LETARTE-MUIRHEAD, M. *see also* BARCLAY, A. N. 699–706
LLEWELLIN, J. M. & GREEN, M. L. The effect of acid proteinase inhibitors on chicken pepsin 319–326
LODER, B. *see* USHER, J. J. 729–739
LOHMANDER, S. *see* THYBERG, J. 157–166

LORD, J. M. Evidence that phosphatidylcholine and phosphatidylethanolamine are synthesized by a single enzyme present in the endoplasmic reticulum of castor-bean endosperm 451–453

MALMSTRÖM, A., CARLSTEDT, I., ÅBERG, L. & FRANSSON, L.-Å. The copolymeric structure of dermatan sulphate produced by cultured human fibroblasts. Different distribution of iduronic acid- and glucuronic acid-containing units in soluble and cell-associated glycans 477–489

MALTHOUSE, J. P. G. *see* STUCHBURY, T. 417–432

MARGISON, G. P., O'CONNOR, P. J. & CORNISH-BOWDEN, A. Role of apurinic sites in the resistance of methylated oligodeoxyribonucleotides to degradation by spleen exonuclease 249–256

MARKEY, F. & WILD, D. G. A 30S precursor of 30S ribosomes in a mutant of *Escherichia coli* 463–465

MARSH, A. *see* ENGELBORGHS, Y. 47–50

MASTERS, C. J. *see* BRINKWORTH, R. I. 631–636; DON, M. M. 625–630

MAYER, R. J. *see* BETTS, S. A. 263–270

MCINTYRE, R. J. *see* VAUGHAN, P. F. T. 759–762

MCLENNAN, A. G. & KEIR, H. M. Deoxyribonucleic acid polymerases of *Euglena gracilis*. Primer-template utilization of and enzyme activities associated with the two deoxyribonucleic acid polymerases of high molecular weight 239–247

MCLENNAN, A. G. & KEIR, H. M. Deoxyribonucleic acid polymerases of *Euglena gracilis*. Purification and properties of two distinct deoxyribonucleic acid polymerases of high molecular weight 227–238

MCNABB, T. & JELLINCK, P. H. Purification and properties of oestrogen-induced uterine peroxidase 275–279

MENON, K. M. J. *see* AZHAR, S. 23–36

MESSER, M. & DEAN, R. T. Immunochemical relationship between α-amylases of rat liver, serum, pancreas and parotid gland 17–22

MEYER, K. *see* CHOI, H. U. 543–553

MICHELACCI, Y. M. & DIETRICH, C. P. A comparative study between a chondroitinase B and a chondroitinase AC from *Flavobacterium heparinum*. Isolation of a chondroitinase AC-susceptible dodecasaccharide from chondroitin sulphate B 121–129

MILSTEIN, C. P., RICHARDSON, N. E., DEVERSON, E. V. & FEINSTEIN, A. Interchain disulphide bridges of mouse immunoglobulin M 615–624

MINATOGAWA, Y. *see* NOGUCHI, T. 399–406

MOLE, L. E. A genetic marker in the variable region of rabbit immunoglobulin heavy chain 351–359

MOLE, L. E. *see also* PRATT, D. M. 337–349; STEVENSON, F. K. 751–753

MORIMOTO, M. *see* NOGUCHI, T. 399–406

MUIR, H. *see* EYRE, D. R. 595–602

MULDER, G. J. & VAN DOORN, A. B. D. A rapid NAD^+-linked assay for microsomal uridine diphosphate glucuronyltransferase of rat liver and some observations on substrate specificity of the enzyme 131–140

MULLINGER, R. N., CAMMACK, R., RAO, K. K., HALL, D. O., DICKSON, D. P. E., JOHNSON, C. E., RUSH, J. D. & SIMOPOULOS, A. Physicochemical characterization of the four-iron–four-sulphide ferredoxin from *Bacillus stearothermophilus* 75–83

NAKATANI, M. *see* NOGUCHI, T. 399–406

NELSON, N. F. *see* BROWN, K. B. 505–512

NOGUCHI, T., MINATOGAWA, Y., OKUNO, E., NAKATANI, M., MORIMOTO, M. & KIDO, R. Purification and characterization of kynurenine–2-oxoglutarate aminotransferase from the liver, brain and small intestine of rats 399–406

NORRIS, R. *see* STUCHBURY, T. 417–432

NORTHCOTE, D. H. *see* YUNG, K.-H. 141–144

NUNEZ, E. *see* LAURENT, C. 513–518

O'CONNOR, P. J. *see* MARGISON, G. P. 249–256

OFFORD, R. E. *see* HALBAN, P. A. 219–225

OKRETIC, M. C. *see* LARA, F. J. S. 575–580

OKUNO, E. *see* NOGUCHI, T. 399–406

O'MALLEY, B. W. *see* KRALL, J. F. 497–503

OSAKI, S., SEXTON, R. C., PASCUAL, E. & FRIEDEN, E. Iron oxidation and transferrin formation by phosvitin 519–525

OSAKI, S. *see also* DE LEY, M. 561–566

OSBORNE, H. H. & HOLLAWAY, M. R. The investigation of substrate-induced changes in subunit interactions in glyceraldehyde 3-phosphate dehydrogenases by measurement of the kinetics and thermodynamics of subunit exchange 37–45

OTTOLENGHI, P. The reversible delipidation of a solubilized sodium-plus-potassium ion-dependent adenosine triphosphatase from the salt gland of the spiny dogfish 61–66

OVERDIJK, B., VAN DER KROEF, W. M. J., VELTKAMP, W. A. & HOOGHWINKEL, G. J. M. The separation of bovine brain β-*N*-acetyl-D-hexosaminidases. Abnormal gel-filtration behaviour of β-*N*-acetyl-D-glucosaminidase C 257–261

PACE, M. *see* BRENNA, O. 291–296

PARR, S. R., WILSON, M. T. & GREENWOOD, C. The reaction of *Pseudomonas aeruginosa* cytochrome *c* oxidase with carbon monoxide 51–59

PARR, S. R. *see also* BRUNORI, M. 185–188

PASCUAL, E. *see* OSAKI, S. 519–525

PERNIGOTTI, L. *see* BACCINO, F. M. 567–573

PERRELLA, M. *see* BRENNA, O. 291–296

PERRY, S. V. *see* FREARSON, N. 99–107; JACKSON, P. 85–97

PETERSON, J. S. *see* KANESHIRO, C. M. 433–434

PHILLIPS, N., ROBINSON, D. & WINCHESTER, B. Immunological characterization of human liver α-D-mannosidase 469–475

PIETTA, P. G. *see* BRENNA, O. 291–296

PÖSÖ, H., SINERVIRTA, R. & JÄNNE, J. *S*-adenosylmethionine decarboxylase from baker's yeast 67–73

POWELL, D. A., DUCKWORTH, M. & BADDILEY, J. A membrane-associated lipomannan in micrococci 387–397

PRATT, D. M. & MOLE, L. E. Sequence studies on the constant region of the Fd sections of rabbit immunoglobulin G of different allotype 337–349

RAO, K. K. *see* MULLINGER, R. N. 75–83

RAYMONT, C. M. *see* STEVENSON, F. K. 751–753

REITHEL, F. J. *see* KANESHIRO, C. M. 433–434

REYNOLDS, C. D. *see* GOODGAME, D. M. L. 467–468

REYNOLDS, J. J. *see* WERB, Z. 645–653, 655–663, 665–669

RICHARDS, R. *see* HARWOOD, J. L. 707–714

RICHARDS, R. E. *see* SIM, E. 555–560

RICHARDSON, N. E. *see* MILSTEIN, C. P. 615–624

ROBINSON, D. *see* PHILLIPS, N. 469–475
ROBINSON, H. C. *see* HOPWOOD, J. J. 581–594
RUSH, J. D. *see* MULLINGER, R. N. 75–83

SAMARAJEEWA, P. & KELLIE, A. E. The radioimmunoassay of steroid glucuronides. The oestrogen C-3 glucuronides as haptens 369–376
SEXTON, R. C. *see* OSAKI, S. 519–525
SHIPTON, M. *see* STUCHBURY, T. 417–432
SIDHU, R. S. & BLAIR, A. H. The action of chelating agents on human liver aldehyde dehydrogenase 443–445
SIM, E., CULLIS, P. R. & RICHARDS, R. E. Physical studies on phosphonium phosphatidylcholine. A unique[^{31}P]-phosphorus nuclear-magnetic-resonance probe for model and biological membranes 555–560
SIMMONDS, R. J. *see* YON, R. J. 281–290
SIMOPOULOS, A. *see* MULLINGER, R. N. 75–83
SINERVIRTA, R. *see* PÖSÖ, H. 67–73
SKAPSKI, A. C. *see* GOODGAME, D. M. L. 467–468
SOCHER, S. H. *see* KRALL, J. F. 497–503
STEINERT, P. M. & IDLER, W. W. The polypeptide composition of bovine epidermal α-keratin 603–614
STEVENSON, F. K., MOLE, L. E., RAYMONT, C. M. & STEVENSON, G. T. A λ Bence-Jones protein in guinea pigs 751–753
STEVENSON, G. T. *see* STEVENSON, F. K. 751–753
STEVENSON, K. J. & GAUCHER, G. M. The substrate specificity of thermomycolase, an extracellular serine proteinase from the thermophilic fungus *Malbranchea pulchella* var. *sulfurea* 527–542
STORER, A. C., DARLISON, M. G. & CORNISH-BOWDEN, A. The nature of experimental error in enzyme kinetic measurements 361–367
STUCHBURY, T., SHIPTON, M., NORRIS R., MALTHOUSE, J. P. G., BROCKLEHURST, K., HERBERT, J. A. L. & SUSCHITZKY, H. A reporter group delivery system with both absolute and selective specificity for thiol groups and an improved fluorescent probe containing the 7-nitrobenzo-2-oxa-1,3-diazole moiety 417–432
SUSCHITZKY, H. *see* STUCHBURY, T. 417–432

TAUB, F. & JOHNSON, T. C. The mechanism of polyribosome disaggregation in brain tissue by phenylalanine 173–180
TETLEY, T. *see* HARWOOD, J. L. 707–714
THIMMAPPAYA, B. & CHERAYIL, J. D. Isoaccepting lysine transfer ribonucleic acid species of *Pseudomonas aeruginosa* 377–386
THOMAS, P. *see* EDWARDS, R. G. 145–148
THYBERG, J., LOHMANDER, S. & HEINEGÅRD, D. Proteoglycans of hyaline cartilage. Electron-microscopic studies on isolated molecules 157–166

USHER, J. J., LODER, B. & ABRAHAM, E. P. Synthesis of tritium-labelled isopenicillin N, penicillin N and 6-aminopenicillanic acid 729–739
USHER, J. J. *see also* FAWCETT, P. A. 741–746

VAN, N. T. *see* KRALL, J. F. 497–503
VAN DER KROEF, W. M. J. *see* OVERDIJK, B. 257–261
VAN DOORN, A. B. D. *see* MULDER, G. J. 131–140
VAUGHAN, P. F. T. & MCINTYRE, R. J. The action of hydrogen peroxide on the hydroxylation of *p*-coumaric acid by spinach-beet phenolase 759–762
VELTKAMP, W. A. *see* OVERDIJK, B. 257–261
VERBUGGEN, R. Subtilopeptidase A isoenzyme system. Interaction with serum components and its importance for quantitative immunoelectrophoresis 149–155
VLIEGENTHART, J. F. G. *see* KAMERLING, J. P. 491–495

WAEHNELDT, T. V. Ontogenetic study of a myelin-derived fraction with 2′:3′-cyclic nucleotide 3′-phosphohydrolase activity higher than that of myelin 435–437
WALKER, J. A. & WHEELER, K. P. Differential effects of temperature on a membrane adenosine triphosphatase and associated phosphatase 439–442
WARD, P. E., GEDNEY, C. D., DOWBEN, R. M. & ERDÖS, E. G. Isolation of membrane-bound renal kallikrein and kininase 755–758
WEIDEMANN, M. J. *see* HICKMAN, P. E. 327–336
WEITZMAN, P. D. J. *see* HARFORD, S. 455–458
WERB, Z. & REYNOLDS, J. J. Immunochemical studies with a specific antiserum to rabbit fibroblast collagenase 655–663
WERB, Z. & REYNOLDS, J. J. Purification and properties of a specific collagenase from rabbit synovial fibroblasts 645–653
WERB Z. & REYNOLDS, J. J. Rabbit collagenase. Immunological identity of the enzymes released from cells and tissues in normal and pathological conditions. 665–669
WESTWOOD, J. H. *see* EDWARDS, R. G. 145–148
WHEELER, K. P. *see* WALKER, J. A. 439–442
WHITTENBURY, R. *see* COLBY, J. 459–462
WILD, D. G. *see* MARKEY, F. 463–465
WILLIAMS, A. F. *see* LETARTE-MUIRHEAD, M. 685–697; 699–706
WILSON, M. T. *see* BRUNORI, M. 185–188; PARR, S. R. 51–59
WINCHESTER, B. *see* PHILLIPS, N. 469–475
WINZOR, D. J. *see* BRINKWORTH, R. I. 631–636; DON, M. M. 625–630

YAMAGUCHI, I. *Oplophorus* oxyluciferin and a model luciferin compound biologically active with *Oplophorus* luciferase 9–15
YON, R. J. & SIMMONDS, R. J. Protein chromatography on adsorbents with hydrophobic and ionic groups. Some properties of *N*-(3-carboxypropionyl)aminodecyl-Sepharose and its interaction with wheat-germ aspartate transcarbamoylase 281–290
YUNG, K.-H. & NORTHCOTE, D. H. Some enzymes present in the walls of mesophyll cells of tobacco leaves 141–144

ZURETTI, M. F. *see* BACCINO, F. M. 567–573

Index of Subjects

β-*N*-Acetylgalactosaminidase, purification and properties of β-*N*-acetylhexosaminidase with activity of β-*N*-acetylglucosaminidase and, from *Trichomonas foetus* (Edwards, R. G., Thomas, P. & Westwood, J. H.) 145–148

β-*N*-Acetyl-D-glucosaminidase C, cerebral-cortex, ox, abnormal gel-filtration behaviour of (Overdijk, B., van der Kroef, W. M. J., Veltkamp, W. A. & Hooghwinkel, G. J. M.) 257–261

β-*N*-Acetylglucosaminidase, purification and properties of β-*N*-acetylhexosaminidase with activity of β-*N*-acetylgalactosaminidase and, from *Trichomonas foetus* (Edwards, R. G., Thomas, P. & Westwood, J. H.) 145–148

β-*N*-Acetylhexosaminidase, purification and properties of, from *Trichomonas foetus* (Edwards, R. G., Thomas, P. & Westwood, J. H.) 145–148

β-*N*-Acetyl-D-hexosaminidases, cerebral-cortex, ox, abnormal gel-filtration behaviour of the β-*N*-acetyl-D-glucosaminidase C component of (Overdijk, B., van der Kroef, W. M. J., Veltkamp, W. A. & Hooghwinkel, G. J. M.) 257–261

Acid α-D-mannosidase, *see* α-D-Mannosidase, acid

Acid phosphatase, *see* Phosphatase, acid

Acid proteinase, *see* Proteinase, acid

α-Actinin, purification of myosin, tropomyosin and, from water-bug and dung-beetle flight muscle (Hammond, K. S. & Goll, D. E.) 189–192

Adenosine triphosphatase, sodium-plus-potassium ion-dependent, membrane, microsomal, kidney, rabbit, differential effects of temperature on the activity of, and on the associated phosphatase activity (Walker, J. A. & Wheeler, K. P.) 439–442

Adenosine triphosphatase, sodium-plus-potassium ion-dependent, microsomal, salt-gland, dogfish, solubilized, reversible delipidation of (Ottolenghi, P.) 61–66

S-Adenosylmethionine decarboxylase, purification and properties of, from baker's yeast (Pösö, H., Sinervirta, R. & Jänne, J.) 67–73

Alanine ester residues, influence of, on the binding of magnesium ions to teichoic acids from *Staphylococcus aureus* H cell wall (Lambert, P. A., Hancock, I. C. & Baddiley, J.) 671–676

Albumin, plasma, capuchin-monkey, role of, in the transport of 25-hydroxycholecalciferol (Hay, A. W. M.) 193–196

Albumin, serum, capuchin monkey, role of, in the transport of 25-hydroxycholecalciferol (Hay, A. W. M.) 193–196

Albumin, serum, human, isolation and properties of a fragment of, demonstrating the absence of a methionine residue from some of the molecules (Lapresle, C. & Doyen, N.) 637–643

Albumin, serum, ox, absolute and selective specificity of the reactions of the fluorescent compounds 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole and 4-(*N*-2-aminoethyl 2′-pyridyl disulphide)-7-nitrobenzo-2-oxa-1,3-diazole with thiol groups of, and other proteins (Stuchbury, T., Shipton, M., Norris, R., Malthouse, J. P. G., Brocklehurst, K., Herbert, J. A. L. & Suschitzky, H.) 417–432

Albumin, serum, ox, theoretical approach to the binding of amphipathic molecules to, and other globular proteins (Jones, M. N.) 109–114

Aldehyde dehydrogenase, liver, human, inhibition by chelating agents of the activity of (Sidhu, R. S. & Blair, A. H.) 443–445

Aldehyde dehydrogenases, cytoplasmic and mitochondrial, liver, sheep, effects of disulfiram on the activities of (Kitson, T. M.) 407–412

Aldolase, *see* Fructose 1,6-diphosphate aldolase

Alkaline phosphatase, *see* Phosphatase, alkaline

Allotypes Aa1 and Aa3, amino acid sequence of the constant region of Fd fragments of the heavy chain of rabbit immunoglobulin G of (Pratt, D. M. & Mole, L. E.) 337–349

Allotypes Aa1 and Aa3, demonstration of a genetic marker in the amino acid sequence of the variable region of the heavy chain of rabbit immunoglobulin G of (Mole, L. E.) 351–359

Amino acid compositions of different molecular species of collagen from pig fibrous, elastic and hyaline cartilages (Eyre, D. R. & Muir, H.) 595–602

Amino acid sequence of *Staphylococcus aureus* N.C.I.B. 11195 penicillinase (Ambler, R. P.) 197–218

Amino acid sequence of the constant region of Fd fragments of the heavy chain of rabbit immunoglobulin G of allotypes Aa1 and Aa3 (Pratt, D. M. & Mole, L. E.) 337–349

Amino acid sequence of the variable region of the heavy chain of rabbit immunoglobulin G of allotypes Aa1 and Aa3, demonstration of a genetic marker in (Mole, L. E.) 351–359

4-(*N*-2-Aminoethyl 2′-pyridyl disulphide)-7-nitrobenzo-2-oxa-1,3-diazole, absolute and selective specificity of the reactions of the fluorescent compounds 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole and, with thiol groups of proteins (Stuchbury, T., Shipton, M., Norris, R., Malthouse, J. P. G., Brocklehurst, K., Herbert, J. A. L. & Suschitzky, H.) 417–432

6-Aminopenicillanic acid, behaviour of tritium-labelled isopenicillin N and, as potential penicillin precursors in an extract of *Penicillium chrysogenum* (Fawcett, P. A., Usher, J. J. & Abraham, E. P.) 741–746

6-Aminopenicillanic acid, synthesis of tritium-labelled isopenicillin N, penicillin N and (Usher, J. J., Loder, B. & Abraham, E. P.) 729–739

α-Amylases, immunochemical relationship between, of rat liver, serum, pancreas and parotid gland (Messer, M. & Dean, R. T.) 17–22

Angiotensin I-converting enzyme, membrane-bound, kidney-cortex, rat, isolation of, and kallikrein (Ward, P. E., Gedney, C. D., Dowden, R. M. & Erdös, E. G.) 755–758

Antigens, use of, in the characterization of Thy-1 glycoprotein purified from rat brain membranes (Barclay, A. N., Letarte-Muirhead, M. & Williams, A. F.) 699–706

Antigens, use of, in the characterization of Thy-1 glycoprotein purified from rat thymocyte membranes (Letarte-Muirhead, M., Barclay, A. N. & Williams, A. F.) 685–697

Anti-(oestradiol-6-carboxymethyloxime–bovine serum albumin) serum, comparison of the oestradiol-17β-binding specificities of rat α-foetoprotein, rat liver 17β-hydroxy steroid dehydrogenase and (Laurent, C., de Lauzon, S., Cittanova, N., Nunez, E. & Jayle, M.-F.) 513–518

Anti-(rabbit synovial-fibroblast collagenase) serum, specific, immunochemical studies with (Werb, Z. & Reynolds, J. J.) 655–663
Apotransferrin, oxidation of ferrous iron and the formation of transferrin from, catalysed by chicken egg-yolk phosvitin (Osaki, S., Sexton, R. C., Pascual, E. & Frieden, E.) 519–525
Arylesterase, plasma, ox, further evidence for the lipoprotein nature of (Don, M. M., Masters, C. J. & Winzor, D. J.) 625–630
Aspartate, activation by, of pig spleen phosphofructokinase (Hickman, P. E. & Weidemann, M. J.) 327–336
Aspartate carbamoyltransferase, wheat-germ, interaction of *N*-(3-carboxypropionyl)aminodecyl-Sepharose with (Yon, R. J. & Simmonds, R. J.) 281–290
Aspartate, transamination of, by rat liver, brain and small-intestine kynurenine–2-oxoglutarate aminotransferase (Noguchi, T., Minatogawa, Y., Okuno, E., Nakatani, M., Morimoto, M. & Kido, R.) 399–406
Azurin, *Pseudomonas aeruginosa* N.C.T.C. 6750, temperature-jump study of the reaction between, and cytochrome *c*-551 oxidase (Brunori, M., Parr, S. R., Greenwood, C. & Wilson, M. T.) 185–188

Bacillus megaterium, evidence from multiple-inhibition studies for isosteric and allosteric inhibition by nucleotides of the activity of citrate synthases from, and other sources (Harford, S. & Weitzman, P. D. J.) 455–458
Bacillus stearothermophilus B65, characterization of poly(glucosylglycerol phosphate) teichoic acid in the cell wall of (Anderson, A. J. & Archibald, A. R.) 115–120
Bacillus stearothermophilus, binding of oxidized nicotinamide-adenine dinucleotide to glyceraldehyde 3-phosphate dehydrogenase from (Allen, G. & Harris, J. I.) 747–749
Bacillus stearothermophilus, physicochemical characterization of the four-iron–four-sulphide ferredoxin from (Mullinger, R. N., Cammack, R., Rao, K. K., Hall, D. O., Dickson, D. P. E., Johnson, C. E., Rush, J. D. & Simopoulos, A.) 75–83
Bean, castor, (*Ricinus communis*), evidence that the biosynthesis of phosphatidylcholine and phosphatidylethanolamine is catalysed by a single enzyme in the endoplasmic reticulum of the endosperm of (Lord, J. M.) 451–453
Beet (*Beta vulgaris*), action of hydrogen peroxide on the hydroxylation of *p*-coumarate by phenolase from (Vaughan, P. F. T. & McIntyre, R. J.) 759–762
Bence-Jones protein, λ, identification of, in the urine of guinea pigs with L_2C lymphocytic leukaemia (Stevenson, F. K., Mole, L. E., Raymont, C. M. & Stevenson, G. T.) 751–753
Beta vulgaris, *see* Beet
Bioluminescent shrimp, *see* Shrimp, bioluminescent
Brain cortex, ox, abnormal gel-filtration behaviour of β-*N*-acetyl-D-glucosaminidase C from (Overdijk, B., van der Kroef, W. M. J., Veltkamp, W. A. & Hooghwinkel, G. J. M.) 257–261
Brain, mouse, neonatal, mechanism of the disaggregation by phenylalanine of polyribosomes in (Taub, F. & Johnson, T. C.) 173–179
Brain, rat, ontogenic study of a myelin-derived membrane fraction with higher activity of 2′:3′-cyclic nucleotide 3′-phosphohydrolase than that of the parent myelin from (Waehneldt, T. V.) 435–437
Brain, rat, purification and characterization of kynurenine-2-oxoglutarate aminotransferases from, and other tissues (Noguchi, T., Minatogawa, Y., Okuno, E., Nakatani, M., Morimoto, M. & Kido, R.) 399–406
Brain, rat, purification and characterization of Thyl-1 glycoprotein from membranes of (Barclay, A. N., Letarte-Muirhead, M. & Williams, A. F.) 699–706
Bromomethane, oxidation of, by *Methylomonas methanica* methane mono-oxygenase (Colby, J., Dalton, H. & Whittenbury, R.) 459–462

Cadmium derivative, X-ray determination of the structure of, of cytidine 5′-monophosphate (Goodgame, D. M. L., Jeeves, I., Reynolds, C. D. & Skapski, A. C.) 467–468
Caeruloplasmin, *see* Ferroxidase
Capuchin monkey, *see* Monkey, capuchin
Carbon monoxide, reaction of, with *Pseudomonas aeruginosa* N.C.T.C. 6750 cytochrome *c*-551 oxidase (Parr, S. R., Wilson, M. T. & Greenwood, C.) 51–59
N-(3-Carboxypropionyl)aminodecyl-Sepharose, properties of, and its interaction with wheat-germ aspartate carbamoyltransferase (Yon, R. J. & Simmonds, R. J.) 281–290
Cardiac muscle, *see* Muscle, heart
Carotenoids, nomeclature of (IUPAC Commission on the Nomenclature of Organic Chemistry & IUPAC–IUB Commission on Biochemical Nomeclature) 5–7
Cartilage, costal, human, structure of keratan sulphates from, and other sources (Choi, H. U. & Meyer, K.) 543–553
Cartilage, hyaline, guinea-pig and ox, electron-microscopic studies on proteoglucans isolated from (Thyberg, J., Lohmander, S. & Heinegård, D.) 157–166
Cartilage, nasal-septum, ox, identification of polydispersity of the chondroitin sulphate component in three proteoglycans from (Hopwood, J. J. & Robinson, H. C.) 581–594
Cartilages, fibrous, elastic and hyaline, pig, distribution of different molecular species of collagen in (Eyre, D. R. & Muir, H.) 595–602
Castor bean, *see* Bean, castor
Cebus albifrons, *see* Monkey, capuchin
Cell wall, *Bacillus stearothermophilus* B65, characterization of poly(glucosylglycerol phosphate) teichoic acid in (Anderson, A. J. & Archibald, A. R.) 115–120
Cell wall, mesophyll-cell, tobacco-leaf, activities and other properties of enzymes in (Yung, K.-H. & Northcote, D. H.) 141–144
Cell wall, *Staphylococcus aureus* H, influence of alanine ester residues on the binding of magnesium ions to teichoic acids from (Lamberg, P. A., Hancock, I. C. & Baddiley, J.) 671–676
Cells, fibroblast, human, cultured, differences in the distribution of units containing iduronic acid and glucuronic acid in soluble and cell-associated dermatan sulphate–chondroitin sulphate co-polymers produced by (Malmström, A., Carlstedt, I., Åberg, L. & Fransson, L.-Å) 477–489
Cells, synovial-fibroblast, rabbit, cultured, immunochemical studies with a specific antiserum to collagenase from (Werb, Z. & Reynolds, J. J.) 655–663
Cells, synovial-fibroblast, rabbit, cultured, immunological identity of the collagenase from, with the enzymes released from other rabbit cells and tissues (Werb, Z. & Reynolds, J. J.) 665–669

Cells, synovial-fibroblast, rabbit, cultured, purification and properties of a specific collagenase from (Werb, Z. & Reynolds, J. J.) 645–653
Cerebellum, mouse, neonatal, mechanism of the disaggregation by phenylalanine of polyribosomes in, and cerebral cortex (Taub, F. & Johnson, T. C.) 173–179
Cerebral cortex, mouse, neonatal, mechanism of the disaggregation by phenylalanine of polyribosomes in, and cerebellum (Taub, F. & Johnson, T. C.) 173–179
Cerebral cortex, ox, abnormal gel-filtration behaviour of *β*-*N*-acetyl-D-glucosaminidase C from (Overdijk, B., van der Kroef, W. M. J., Veltkamp, W. A. & Hooghwinkel, G. J. M.) 257–261
Chick, resistance to nuclease of chromatin deoxyribonucleic acid from the oviduct of (Krall, J. F., Socher, S. H., Van, N. T. & O'Malley, B. W.) 497–503
Chicken, crystallization of one of the pepsinogens from, and of the derived pepsin (Green, M. L.) 763–764
Chicken, effects of inhibitors of acid proteinases on the activity of pepsin from (Llewellin, J. M. & Green, M. L.) 319–326
Chicken egg, oxidation of ferrous iron and the formation of transferrin from apotransferrin catalysed by egg-yolk phosvitin from (Osaki, S., Sexton, R. C., Pascual, E. & Frieden, E.) 519–525
4-Chloro-7-nitrobenzo-2-oxa-1,3-diazole, absolute and selective specificity of the reactions of the fluorescent compounds 4-(*N*-2-aminoethyl 2′-pyridyl disulphide)-7-nitrobenzo-2-oxa-1,3-diazole and, with thiol groups of proteins (Stuchbury, T., Shipton, M., Norris, R., Malthouse, J. P. G., Brocklehurst, K., Herbert, J. A. L. & Suschitzky, H.) 417–432
Cholecalciferol, 25-hydroxy-, transport of, in the plasma of the capuchin monkey (Hay, A. W. M.) 193–196
Choline dehydrogenase, mitochondrial, liver, rat, reaction of, with some electron acceptors (Barrett, M. C. & Dawson, A. P.) 677–683
Chondroitin sulphate B, *Flavobacterium heparinum* A.T.C.C. 13125, isolation of a dodecasaccharide from, that is susceptible to chondroitinase AC from the same species (Michelacci, Y. M. & Dietrich, C. P.) 121–129
Chondroitin sulphate, differences in the distribution of units containing iduronic acid and glucuronic acid in soluble and cell-associated co-polymers of dermatan sulphate and, produced by cultured human fibroblasts (Malmström, A., Carlstedt, I., Åberg, L. & Fransson, L.-Å.) 477–489
Chondroitin sulphate, identification of polydispersity in, in three ox nasal-septum cartilage proteoglycans (Hopwood, J. J. & Robinson, H. C.) 581–594
Chondroitinase AC, *Flavobacterium heparinum* A.T.C.C. 13125, isolation of a dodecasaccharide from chondroitin B that is susceptible to (Michelacci, Y. M. & Dietrich, C. P.) 121–129
Chromatin, oviduct, chick, resistance to nuclease of deoxyribonucleic acid of (Krall, J. F., Socher, S. H., Van, N. T. & O'Malley, B. W.) 497–503
Chymosin, calf, inhibition by 1,2-epoxy-3-(*p*-nitrophenoxy)propane of the activities of chicken pepsin and (Llewellin, J. M. & Green, M. L.) 319–326
α-Chymotrypsin, ox, nature of experimental error in kinetic measurements of reactions catalysed by, and other enzymes (Storer, A. C., Darlison, M. G. & Cornish-Bowden, A.) 361–367
Citrate synthases, evidence from multiple-inhibition studies for isosteric and allosteric inhibition by nucleotides of the activity of, from pig heart, *Bacillus megaterium* and *Pseudomonas aeruginosa* (Harford, S. & Weitzman, P. D. J.) 455–458
Clostridium perfringens, use of affinity chromatography in an improved method for the purification of sialidase from (Geosow, M. J.) 181–183
Colchicine, radioimmunoassay of (Boudene, C., Duprey, F. & Bohuon, C.) 413–415
Collagen, distribution of different molecular species of, in pig fibrous, elastic and hyaline cartilages (Eyre, D. R. & Muir, H.) 595–602
Collagenase, specific, purification and properties of, from cultured rabbit synovial fibroblasts (Werb, Z. & Reynolds, J. J.) 645–653
Collagenase, synovial-fibroblast, rabbit, immunochemical studies with a specific antiserum to (Werb, Z. & Reynolds, J. J.) 655–663
Collagenase, synovial-fibroblast, rabbit, immunological identity of, with the enzymes released from other rabbit cells and tissues (Werb, Z. & Reynolds, J. J.) 665–669
Cornea, ox, structure of keratan sulphates from, and other sources (Choi, H. U. & Meyer, K.) 543–553
Corpus luteum, ox, properties of adenosine 3′:5′-cyclic monophosphate-dependent protein kinases associated with the plasma membrane of (Azhar, S. & Menon, K. M. J.) 23–36
p-Coumarate, action of hydrogen peroxide on the hydroxylation of, by spinach-beet phenolase (Vaughan, P. F. T. & McIntyre, R. J.) 759–762
Creatine kinase, skeletal-muscle, rabbit, quenched-flow study of the reaction catalysed by (Engelborghs, Y., Marsh, A. & Gutfreund, H.) 47–50
2′:3′-Cyclic nucleotide phosphohydrolase, ontogenic study of a myelin-derived membrane fraction with higher activity of, than that of the parent myelin from rat forebrain (Waehneldt, T. V.) 435–437
Cytidine 5′-monophosphate, X-ray determination of the structure of a cadmium derivative of (Goodgame, D. M. L., Jeeves, I., Reynolds, C. D. & Skapski, A. C.) 467–468
Cytochrome *c*-551 oxidase, *Pseudomonas aeruginosa* N.C.T.C. 6750, reaction of carbon monoxide with (Parr, S. R., Wilson, M. T. & Greenwood, C.) 51–59
Cytochrome *c*-551 oxidase, *Pseudomonas aeruginosa* N.C.T.C. 6750, temperature-jump study of the reaction between, and azurin (Brunori, M., Parr, S. R., Greenwood, C. & Wilson, M. T.) 185–188
Cytosol, liver, sheep, effects of disulfiram on the activities of aldehyde dehydrogenases from, and mitochondria (Kitson, T. M.) 407–412

Deoxyribonuclease, exo-, spleen, pig, role of apurinic sites in the resistance of methylated oligodeoxyribonucleotides to degradation by (Margison, G. P., O'Connor, P. J. & Cornish-Bowden, A.) 249–256
Deoxyribonucleic acid, chromatin, oviduct, chick, resistance to nuclease of (Krall, J. F., Socher, S. H., Van N. T. & O'Malley, B. W.) 497–503
Deoxyribonucleic acid, nuclear, liver, rat, effects of polyamines and methylglyoxal bis(guanylhydrazone) on the template activity of (Brown, K. B., Nelson, N. F. & Brown, D. G.) 505–512

Deoxyribonucleic acid polyermerases, high molecular-weight, two, purification and properties of, from *Euglena gracilis* (McLennan, A. G. & Keir, H. M.) 227–238
Deoxyribonucleic acid polymerases, high-molecular-weight, two, utilization of primer template by and enzyme activities associated with, from *Euglena gracilis* (McLennan, A. G. & Keir, H. M.) 239–247
Dermatan sulphate, differences in the distribution of units containing iduronic acid and glucuronic acid in soluble and cell-associated co-polymers of chondroitin sulphate and, produced by cultured human fibroblasts (Malmström, A., Carlstedt, I., Åberg, L. & Fransson, L.-Å.) 477–489
Diazoacetylnorleucine methyl ester, effects of, and other inhibitors of acid proteinases on the activity of chicken pepsin (Llewellin, J. M. & Green, M. L.) 319–326
2-Diazo-4-bromoacetophenone, effects of, and other inhibitors of acid proteinases on the activity of chicken pepsin (Llewellin, J. M. & Green, M. L.) 319–326
Dimethyl-(2-hydroxy-5-nitrobenzyl)sulphonium bromide, effects of, and other inhibitors of acid proteinases on the activity of chicken pepsin (Llewellin, J. M. & Green, M. L.) 319–326
Disulfiram (tetraethylthiuram disulphide), effects of, on the activities of sheep liver cytoplasmic and mitochondrial aldehyde dehydrogenases (Kitson, T. M.) 407–412
Disulphide bridges, interchain, arrangement of, of mouse immunoglobulin M (Milstein, C. P., Richardson, N. E., Deverson, E. V. & Feinstein, A.) 615–624
Dodecasaccharide, isolation of, from *Flavobacterium heparinum* A.T.C.C. 13125 chondroitin sulphate B that is suceptible to chondroitinase AC from the same species (Michelacci, Y. M. & Dietrich, C. P.) 121–129
Dodecyl sulphate, sodium, theoretical approach to the binding of amphipathic molecules such as, to globular proteins (Jones, M. N.) 109–114
Dogfish (*Squalus acanthias*), reversible delipidation of solubilized microsomal sodium-plus-potassium ion-dependent adenosine triphosphatase from the rectal salt gland of (Ottolenghi, P.) 61–66
Dogfish (*Squalus acanthius*), reversible reaction of pyridoxal 5′-phosphate with an essential lysine residue of skeletal-muscle lactate dehydrogenase M_4 isoenzyme from (Chen, S.-S. & Engel, P. C.) 447–449
Dung beetle (*Heliocopris japetus*), purification of myosin, tropomyosin and α-actinin from the flight muscle of (Hammond, K. S. & Goll, D. E.) 189–192

Egg yolk, chicken, oxidation of ferrous iron and the formation of transferin from apotransferrin catalysed by phosvitin from (Osaki, S., Sexton, R. C., Pascual, E. & Frieden, E.) 519–525
Elastase, activity of, of the extracellular serine proteinase thermomycolin from *Malbranchea pulchella* var. *sulfurea* (Stevenson, K. J. & Gaucher, G. M.) 527–542
Electron transfer, temperature-jump study of, between *Pseudomonas aeruginosa* N.C.T.C. 6750 azurin and cytochrome *c*-551 oxidase (Brunori, M., Parr, S. R., Greenwood, C. & Wilson, M. T.) 185–188
Electron transport, intramolecular, in human ferroxidase (De Ley, M. & Osaki, S.) 561–566
Endoplasmic reticulum, *see* Reticulum, endoplasmic
Endosperm, castor-bean, evidence that the biosynthesis of phosphatidylcholine and phosphatidylethanolamine is catalysed by a single enzyme in the endoplasmic reticulum of (Lord, J. M.) 451–453
Enzymes, nature of experimental error in kinetic measurements of reactions catalysed by (Storer, A. C., Darlison, M. G. & Cornish-Bowden, A.) 361–367
Epidermis, ox, characterization of the polypeptide chains of α-keratin from (Steinert, P. M. & Idler, W. W.) 603–614
1,2-Epoxy-3-(*p*-nitrophenoxy)propane, effects of, and other inhibitors of acid proteinases on the activity of chicken pepsin (Llewellin, J. M. & Green, M. L.) 319–326
Escherichia coli, accumulation of a 30S precursor of 30S ribosomes in a mutant of (Markey, F. & Wild, D. G.) 463–465
Escherichia coli, evidence for an active dimer of β-galactosidase from (Kaneshiro, C. M., Enns, C. A., Hahn, M. G., Peterson, J. S. & Reithel, F. J.) 433–434
Estradiol, *see* Oestradiol
Estradiol-17β, *see* Oestradiol-17β
Estriol, *see* Oestriol
Estrogens, *see* Oestrogens
Estrone, *see* Oestrone
Euglena gracilis, purification and properties of two distinct high-molecular-weight deoxyribonucleic acid polymerases from (McLennan, A. G. & Keir, H. M.) 227–238
Euglena gracilis, utilization of primer template by and enzyme activities associated with two high-molecular-weight deoxyribonucleic acid polymerases from (McLennan, A. G. & Keir, H. M.) 239–247
Exodeoxyribonuclease, spleen, pig, role of apurinic sites in the resistance of methylated oligodeoxyribonucleotides to degradation by (Margison, G. P., O'Connor, P. J. & Cornish-Bowden, A.) 249–256

Ferrodoxin, four-iron–four-sulphide, *Bacillus stearothermophilus*, physicochemical characterization of (Mullinger, R. N., Cammack, R., Rao, K. K., Hall, D. O., Dickson, D. P. E., Johnson, C. E., Rush, J. D. & Simopoulos, A.) 75–83
Ferrous iron, oxidation of, and the formation of transferrin from apotransferrin catalysed by chicken egg-yolk phosvitin (Osaki, S., Sexton, R. C., Pascual, E. & Frieden, E.) 519–525
Ferroxidase, human, intramolecular electron transport in (De Ley, M. & Osaki, S.) 561–566
Ferroxidase, role, of, and phosvitin and transferrin in the mobilization and transfer of iron in the chicken (Osaki, S., Sexton, R. C., Pascaul, E. & Frieden, E.) 519–525
Fibroblasts, human, cultured, differences in the distribution of units containing iduronic acid and glucuronic acid in soluble and cell-associated dermatan sulphate–chondroitin sulphate co-polymers produced by (Malmström, A., Carlstedt, I., Åberg, L. & Fransson, L.-Å.) 477–489
Fibroblasts, synovial, rabbit, cultured, immunochemical studies with a specific antiserum to collagenase from (Werb, Z. & Reynolds, J. J.) 655–663
Fibroblasts, synovial, rabbit, cultured, immunological identity of the collagenase from, with the enzymes released from other rabbit cells and tissues (Werb, Z. & Reynolds, J. J.) 665–669

Fibroblasts, synovial, rabbit, cultured, purification and properties of a specific collagenase from (Werb, Z. & Reynolds, J. J.) 645–653

Ficin, absolute and selective specificity of the reactions of the fluorescent compounds 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole and 4-(*N*-2-aminoethyl 2′-pyridyl disulphide)-7-nitrobenzo-2-oxa-1,3-diazole with thiol groups of, and other proteins (Stuchbury, T., Shipton, M., Norris, R., Malthouse, J. P. G., Brocklehurst, K., Herbert, J. A. L. & Suschitzky, H.) 417–432

Flavobacterium heparinum A.T.C.C. 13125, isolation of a dodecasaccharide from chondroitin sulphate B from, that is susceptible to chondroitinase AC from the same species (Michelacci, Y. M. & Dietrich, C. P.) 121–129

Flight muscle, *see* Muscle, flight

Fly (*Rynchosciara americana*) larva, gel-electrophoretic analysis of salivary-gland messenger ribonucleic acid from (Lara, F. J. S. & Okretic, M. C.) 575–580

α-Foetoprotein, rat, comparison of the oestradiol-17β-binding specificities of rat liver 17β-hydroxy steroid dehydrogenase, anti(oestradiol-6-carboxymethyloxime–bovine serum albumin) serum and (Laurent, C., de Lauzon, S., Cittanova, N., Nunez, E. & Jayle, M.F.) 513–518

Forebrain, rat, ontogenic study of a myelin-derived membrane fraction with higher activity of 2′:3′-cyclic nucleotide 3′-phosphohydrolase than that of the parent myelin from (Waehneldt, T. V.) 435–437

Fructose 1,6-diphosphate aldolase, skeletal-muscle, rabbit, effect of pH on the rate of formation and on the equilibrium concentration of the carbanion intermediate formed in the reaction catalysed by (Grazi, E.) 167–172

β-Galactosidase, *Escherichia coli*, evidence for an active dimer of (Kaneshiro, C. M., Enns, C. A., Hahn, M. G., Peterson, J. S. & Reithel, F. J.) 433–434

β-Galactosidase, liver, rat, stimulation by ions of the activity of (Baccino, F. M., Zuretti, M. F. & Pernigotti, L.) 567–573

Glucokinse, liver, rat, nature of experimental error in kinetic measurements of reactions catalysed by, and other enzymes (Storer, A. C., Darlison, M. G. & Cornish-Bowden, A.) 361–367

Glucuronic acid, differences in the distribution of units containing iduronic acid and, in soluble and cell-associated dermatan sulphate–chondroitin sulphate copolymers produced by cultured human fibroblasts (Malmström, A., Carlstedt, I., Åberg, L. & Fransson, L.-Å.) 477–489

Glutamate dehydrogenase, liver, ox, kinetics of the product inhibition of (Engel, P. C. & Chen, S.-S.) 305–318

Glyceraldehyde 3-phosphate dehydrogenase, *Bacillus stearothermophilus*, binding of oxidized nicotinamide–adenine dinucleotide to (Allen, G & Harris, J. I.) 747–749

Glyceraldehyde 3-phosphate dehydrogenases, baker's-yeast and skeletal-muscle, rabbit, investigation of substrate-induce changes in subunit interactions in, by measurement of the kinetics and thermodynamics of subunit exchange (Osborne, H. H. & Holloway, M. R.) 37–45

Glycopeptides, characterization by gas–liquid chromatography–mass spectrometry and proton-magnetic-resonance spectroscopy of pertrimethylsilyl methyl glycosides obtained in the methanolysis of glycoproteins and (Kamerling, J. P., Gerwig, G. J., Vliegenthart, J. F. G. & Clamp, J. R.) 491–495

Glycoprotein, Thy-1, purification and characterization of, from rat brain membranes (Barclay, A. N., Letarte-Muirhead, M. & Williams, A. F.) 699–706

Glycoprotein, Thy-1, purification and characterization of, from rat thymocyte membranes (Letarte-Muirhead, M., Barclay, A. N. & Williams, A. F.) 685–697

Glycoproteins, characterization by gas–liquid chromatography–mass spectrometry and proton-magnetic-resonance spectroscopy of pertrimethylsilyl methyl glycosides obtained in the methanolysis of glycopeptides and (Kamerling, J. P., Gerwig, G. J., Vliegenthart, J. F. G. and Clamp, J. R.) 491–495

Glycosaminoglycans, soluble and cell-associated, differences in the distribution of units containing iduronic acid and glucuronic acid in, produced by cultured human fibroblasts (Malmström, A., Carlstedt, I., Åberg, L. & Fransson, L.-Å.) 477–489

Glycosaminoglycans, structural variations in, from various sources (Choi, H. U. & Meyer, K.) 543–553

Heart, mouse, evaluation by affinity chromatography of the equilibrium constants for the interaction of reduced nicotinamide–adenine dinucleotide with lactate dehydrogenase isoenzymes from, and other tissues (Brinkworth, R. I., Masters, C. J. & Winzor, D. J.) 631–636

Heart muscle, rabbit, phosphorylation of light-chain components of myosin from, and red skeletal muscle (Frearson, N. & Perry, S. V.) 99–107

Heart, pig, binding of pyruvate to lactate dehydrogenase from, and the interconversion of pyruvate-containing ternary complexes (Boland, M. J. & Gutfreund, H.) 715–727

Heart, pig, evidence from multiple-inhibition studies for isosteric and allosteric inhibition by nucleotides of the activity of citrate synthases from, and other sources (Harford, S. & Weitzman, P. D. J.) 455–458

Heart, pig, reversible modification by pyridoxal 5′-phosphate of mitochondrial malate dehydrogenase from (Chen, S.-S. & Engel, P. C.) 297–303

Heavy chain, immunoglobulin G, rabbit, amino acid sequence of the constant region of Fd fragments of, of allotypes Aa1 and Aa3 (Pratt, D. M. & Mole, L. E.) 337–349

Heavy chain, immunoglobulin G, rabbit, demonstration of a genetic marker in the amino acid sequence of the variable region of, of allotypes Aa1 and Aa3 (Mole, L. E.) 351–359

Heavy chains, immunoglobulin M, mouse, arrangement of the disulphide bridges linking, and light chains (Milstein, C. P., Richardson, N. E., Deverson, E. V. & Feinstein, A.) 615–624

Heliocopris japetus, *see* Dung beetle

Hen, *see* Chicken

Hormones, peptide, nomenclature of (IUPAC–IUB Commission on Biochemical Nomenclature) 1–4

Hyaline cartilage, *see* Cartilage, hyaline

Hydrogen isotope (^3H), semi-synthesis of pig insulin labelled with, with a specific radioactivity of up to 20 curies per millimole (Halban, P. A. & Offord, R. E.) 219–225

Hydrogen isotope (^3H), synthesis of isopenicillin N, penicillin N and 6-aminopenicillanic acid labelled with (Usher, J. J., Loder, B. & Abraham, E. P.) 729–739

Hydrogen peroxide, action of, on the hydroxylation of *p*-coumarate by spinach-beet phenolase (Vaughan, P. F. T. & McIntyre, R. J.) 759–762

17β-Hydroxy steroid dehydrogenase, liver, rat, comparison of the oestradiol-17β-binding specificities of rat α-foetoprotein, anti-(oestradiol-6-carboxymethyloxime–bovine serum albumin) serum and (Laurent, C., de Lauzon, S., Cittanova, N., Nunez, E. & Jayle, M.-F.) 513–518

25-Hydroxycholecalciferol, transport of, in the plasma of the capuchin monkey (Hay, A. W. M.) 193–196

3-Hydroxy-2-methylimidazo[1,2-*a*]pyridine, chemical synthesis of, and its biological activity as a model luciferin compound with luciferase from the bioluminescent shrimp *Oplophorus gracilorostris* (Yamaguchi, I.) 9–15

Iduronic acid, differences in the distribution of units containing glucuronic acid and, in soluble and cell-associated dermatan sulphate–chondroitin sulphate copolymers produced by cultured human fibroblasts (Malmström, A., Carlstedt, I., Åberg, L. & Fransson, L.-Å.) 477–489

Immunoglobulin G, rabbit, amino acid sequence of the constant region of Fd fragments of the heavy chain of, of allotypes Aa1 and Aa3 (Pratt, D. M. & Mole, L. E.) 337–349

Immunoglobulin G, rabbit, demonstration of a genetic marker in the amino acid sequence of the variable region of the heavy chain of, of allotypes Aa1 and Aa3 (Mole, L. E.) 351–359

Immunoglobulin M, mouse, arrangement of the interchain disulphide bridges of (Milstein, C. P., Richardson, N. E., Deverson, E. V. & Feinstein, A.) 615–624

Immunoglobulins, interaction of the subtilopeptidase A isoenzyme system with, and other serum components and its importance for quantitative immunoelectrophoresis (Verbruggen, R.) 149–155

Insulin, pig, ^3H-labelled, semi-synthesis of, with a specific radioactivity of up to 20 curies per millimole (Halban, P. A. & Offord, R. E.) 219–225

Intestine, small, calf, affinity-chromatographic purification of alkaline phosphatase from mucosa of (Brenna, O., Perrella, M., Pace, M. & Pietta, P. G.) 291–296

Intestine, small, rat, purification and characterization of kynurenine–2-oxoglutarate aminotransferases from, and other tissues (Noguchi, T., Minatogawa, Y., Okuno, E., Nakatani, M., Morimoto, M. & Kido, R.) 399–406

Ions, stimulation by, of the activity of rat liver β-galactosidase (Baccino, F. M., Zuretti, M. F. & Pernigotti, L.) 567–573

Iron, ferrous, oxidation of, and the formation of transferrin from apotransferrin catalysed by chicken egg-yolk phosvitin (Osaki, S., Sexton, R. C., Pascual, E. & Frieden, E.) 519–525

Isopenicillin N, behaviour of tritium-labelled 6-aminopenicillanic acid and, as potential penicillin precursors in an extract of *Penicillium chrysogenum* (Fawcett, P. A., Usher, J. J. & Abraham, E. P.) 741–746

Isopenicillin N, synthesis of tritum-labelled 6-aminopenicillanic acid, penicillin N and (Usher, J. J., Loder, B. & Abraham, E. P.) 729–739

Kallikrein, membrane-bound, kidney-cortex, rat, isolation of, and kininase II (Ward, P. E., Gedney, C. D., Dowben, R. M. & Erdös, E. G.) 755–758

Keratan sulphates, structure of, from various sources (Choi, H. U. & Meyer, K.) 543–553

α-Keratin, epidermal, ox, characterization of the polypeptide chains of (Steinert, P. M. & Idler, W. W.) 603–614

Kidney cortex, rat, isolation of membrane-bound kallikrein and kininase II from (Ward, P. E., Gedney, C. D., Dowben, R. M. & Erdös, E. G.) 755–758

Kidney, mouse, evaluation by affinity chromatography of the equilibrium constants for the interaction of reduced nicotinamide–adenine dinucleotide with lactate dehydrogenase isoenzymes from, and other tissues (Brinkworth, R. I., Masters, C. J. & Winzor, D. J.) 631–636

Kidney, rabbit, differential effects of temperature on the activity of membrane sodium-plus-potassium ion-dependent adenosine triphophatase and on the associated phosphatase activity of microsomal fraction from (Walker, J. A. & Wheeler, K. P.) 439–442

Kininase II, membrane-bound, kidney-cortex, rat, isolation of, and kallikrein (Ward, P. E., Gedney, C. D., Dowben, R. M. & Erdös, E. G.) 755–758

Kynurenine–2-oxoglutarate aminotransferases, purification and characterization of, from rat liver, brain and small intestine (Noguchi, T., Minatogawa, Y., Okuno, E., Nakatani, M., Morimoto, M. & Kido, R.) 399–406

β-Lactamase, *Staphylococcus aureus* N.C.I.B. 11195, amino acid sequence of (Ambler, R. P.) 197–218

Lactate dehydrogenase, heart, pig, binding of pyruvate to, and the interconversion of pyruvate-containing ternary complexes (Boland, M. J. & Gutfreund, H.) 715–727

Lactate dehydrogenase, skeletal-muscle, dogfish, reversible reaction of pyridoxal 5′-phosphate with an essential lysine residue of (Chen, S.-S. & Engel, P. C. 447–449

Lactate dehydrogenase, skeletal-muscle, heart, liver and kidney, mouse, evaluation by affinity chromatography of the equilibrium constants for the interaction of isoenzymes of, with reduced nicotinamide–adenine dinucleotide (Brinkworth, R. I., Masters, C. J. & Winzor, D. J.) 631–636

β-Lactoglobulin, theoretical approach to the binding of amphipathic molecules to, and other globular proteins (Jones, M. N.) 109–114

Leaves, tobacco, activities and other properties of enzymes in the cell wall of mesophyll cells of (Yung, K.-H. & Northcote, D. H.) 141–144

Lecithin, *see* Phosphatidylcholine

Lethocerus cordofanus, *see* Water bug

Leucocytes, polymorphonuclear, rabbit, evidence for immunological non-identity of rabbit synovial-fibroblast collagenase with proteinase from (Werb, Z. & Reynolds, J. J.) 665–669

Leukaemia, lymphocytic, L_2C, identification of a λ Bence-Jones protein in the urine of guinea pigs with (Stevenson, F. K., Mole, L. E., Raymont, C. M. & Stevenson, G. T.) 751–753
Ligand, factorization of the Michaelis functions for the binding of, to a molecule with two binding sites (Dixon, H. B. F.) 271–274
Light chains, immunoglobulin M, mouse, arrangement of the disulphide bridges linking, and heavy chains (Milstein, C. P., Richardson, N. E., Deverson, E. V. & Feinstein, A.) 615–624
Light chains, myosin, heart-muscle and red-skeletal-muscle, rabbit, phosphorylation of (Frearson, N. & Perry, S. V.) 99–107
Lipid, content of protein and, in pulmonary surfactants from ox, rabbit, rat and sheep (Harwood, J. L., Desai, R., Hext, P., Tetley, T. & Richards, R.) 707–714
Lipid, reversible removal of, from solubilized dogfish salt gland microsomal sodium-plus-potassium ion-dependent adenosine triphosphatase (Ottolenghi, P.) 61–66
Lipomannans, structure and properties of, associated with membranes of various micrococci (Powell, D. A., Duckworth, M, & Baddiley, J.) 387–397
Lipoprotein, further evidence that ox plasma arylesterase is, in nature (Don, M. M., Masters, C. J. & Winzor, D. J.) 625–630
Lipoteichoic acids, evidence for the absence of, from *Micrococcus lysodeikticus* and other micrococci (Powell, D. A., Duckworth, M. & Baddiley, J.) 387–397
Liver, human, immunological characterization of acid α-D-mannosidases from (Phillips, N., Robinson, D. & Winchester, B.) 469–475
Liver, human, inhibition by chelating agents of the activity of aldehyde dehydrogenase from (Sidhu, R. S. & Blair, A. H.) 443–445
Liver, mouse, evaluation by affinity chromatography of the equilibrium constants for the interaction of reduced nicotinamide–adenine dinucleotide with lactate dehydrogenase isoenzymes from, and other tissues (Brinkworth, R. I., Masters, C. J. & Winzor, D. J.) 631–636
Liver, ox, kinetics of the product inhibition of glutamate dehydrogenase from (Engel, P. C. & Chen, S.-S.) 305–318
Liver, rat, comparison of the oestradiol-17β-binding specificities of rat α-foetoprotein, anti-(oestradiol-6-carboxymethyloxime–bovine serum albumin) serum and 17β-hydroxy steroid dehydrogenase from (Laurent, C., de Lauzon, S., Cittanova, N., Nunez, E. & Jayle, M.-F.) 513–518
Liver, rat, effects of polyamines and methylglyoxal bis-(guanylhydrazone) on nuclear structure and deoxyribonucleic acid template activity in (Brown, K. B., Nelson, N. F. & Brown, D. G.) 505–512
Liver, rat, immunochemical relationship between α-amylases of, and serum, pancreas and parotid gland (Messer, M. & Dean, R. T.) 17–22
Liver, rat, purification and characterization of kynurenine–2-oxoglutarate aminotransferases from, and other tissues (Noguchi, T., Minatogawa, Y., Okuno, E., Nakatani, M., Morimoto, M. & Kido, R.) 399–406
Liver, rat, rapid nicotinamide–adenine dinucleotide-linked method for the assay of the activity of microsomal uridine diphosphate glucuronyltransferase from, and the substrate specificity of the enzyme (Mulder, G. J. & Van Doorn, A. B. D.) 131–140
Liver, rat, reaction of mitochondrial choline dehydrogenase from, with some electron acceptors (Barrett, M. C. & Dawson, A. P.) 677–683
Liver, rat, stimulation by ions of the activity of β-galactosidase in (Baccino, F. M., Zuretti, M. F. & Pernigotti, L.) 567–573
Liver, sheep, effects of disulfiram on the activities of the cytoplasmic and mitochondrial aldehyde dehydrogenases from (Kitson, T. M.) 407–412
Luciferase, bioluminescent-shrimp (*Oplophorus gracilorostris*), activity of, with purified oxyluciferin from the same organism and with a chemically synthesized model luciferin compound (Yamaguchi, I.) 9–15
Luciferin, purification and properties of, from the bioluminescent shrimp *Oplophorus gracilorostris* and the chemical synthesis of a model luciferin compound biologically active with luciferase from the same organism (Yamaguchi, I.) 9–15
Lung, ox, rabbit, rat and sheep, characterization of pulmonary surfactants from (Harwood, J. L., Desai, R., Hext, P., Tetley, T. & Richards, R.) 707–714
Lysine residue, essential, reversible interaction of pyridoxal 5′-phosphate with, of pig heart mitochondrial malate dehydrogenase (Chen, S.-S. & Engel, P. C.) 297–303
Lysine residue, essential, reversible reaction of pyridoxal 5′-phosphate with, of dogfish skeletal-muscle lactate dehydrogenase M_4 isoenzyme (Chen, S.-S. & Engel, P. C.) 447–449
Lysine transfer ribonucleic acid, demonstration of two isoaccepting species of, in *Pseudomonas aeruginosa* (Thimmappaya, B. & Cherayil, J. D.) 377–386
Lysosomes, liver, rat, stimulation by ions of the activity of β-galactosidase in (Baccino, F. M., Zuretti, M. F. & Pernigotti, L.) 567–573

Macroglobulin, *see* Immunoglobulin M
Magnesium chloride, stimulation by and other salts of the activity of rat liver β-galactosidase (Baccino, F. M., Zuretti, M. F. & Pernigotti, L.) 567–573
Magnesium ions, influence of alanine ester residues on the binding of, to teichoic acids from *Staphylococcus aureus* H cell wall (Lambert, P. A., Hancock, I. C. & Baddiley, J.) 671–676
Malate dehydrogenase, mitochondrial, heart, pig, reversible interaction of pyridoxal 5′-phosphate with an essential lysine residue of (Chen, S.-S. & Engel, P. C.) 297–303
Malbranchea pulchella var. *sulfurea*, substrate specificity of the extracellular serine proteinase thermomycolin from (Stevenson, K. J. & Gaucher, G. M.) 527–542
Mammary gland, rabbit, purification and properties of 6-phosphogluconate dehydrogenase from (Betts, S. A. & Mayer, R. J.) 263–270
α-D-Mannosidases, acid, liver, human, immunological characterization of (Phillips, N., Robinson, D. & Winchester, B.) 469–475
Membrane fraction, myelin-derived, forebrain, rat, ontogenic study of, with higher activity of 2′:3′-cyclic nucleotide 3′-phosphohydrolase than that of the parent myelin (Waehneldt, T. V.) 435–437
Membrane, microsomal, kidney, rabbit, differential effects of temperature on the activity of sodium-plus-potassium ion-dependent adenosine triphosphatase and on the associated phosphatase activity of (Walker, J. A. & Wheeler, K. P.) 439–442

Membrane, plasma, corpus luteum, ox, properties of adenosine 3′:5′-cyclic monophosphate-dependent protein kinases associated with (Azhar, S. & Menon, K. M. J.) 23–36
Membranes, brain, rat, purification and characterization of Thy-1 glucoprotein from (Barclay, A. N., Letarte-Muirhead, M. & Williams, A. F.) 699–706
Membranes, kidney-cortex, rat, isolation of kallikrein and kininase II bound to (Ward, P. E., Gedney, C. D., Dowben, R. M. & Erdös, E. G.) 755–758
Membranes, micrococcal, structure and properties of lipomannans associated with (Powell, D. A., Duckworth, M. & Baddiley, J.) 387–397
Membranes, model and biological, physical properties of phosphonium phosphatidylcholine and its suitability as a ^{31}P nuclear-magnetic-resonance probe for (Sim, E. Cullis, P. R. & Richards, R. E.) 555–560
Membranes, thymocyte, rat, purification and characterization of Thy-1 glycoprotein from (Letarte-Muirhead, M., Barclay, A. N. & Williams, A. F.) 685–697
Messenger ribonucleic acid, *see* Ribonucleic acid, messenger
Methane mono-oxygenase, properties of, from *Methylomonas methanica* (Colby, J., Dalton, H. & Whittenbury, R.) 459–462
Methionine residue, isolation and properties of a fragment of human serum albumin demonstrating the absence of, from some of the molecules (Lapresle, C. & Doyen, N.) 637–643
Methyl bromide, *see* Bromomethane
Methyl glycosides, pertrimethylsilyl, characterization by gas–liquid chromatography–mass spectrometry and proton-magnetic-resonance spectroscopy of, obtained in the methanolysis of glycoproteins and glycopeptides (Kamerling, J. P., Gerwig, G. J., Vliegenthart, J. F. G. & Clamp, J. R.) 491–495
Methylglyoxal bis(guanylhydrazone) {1,1′-[(methylethanediylidene)dinitrilo]diguanidine}, effects of polyamines and, on rat liver nuclear structure and deoxyribonucleic acid template activity (Brown, K. B., Nelson, N. F. & Brown, D. G.) 505–512
Methylomonas methanica, properties of methane mono-oxygenase from (Colby, J., Dalton, H. & Whittenbury, R.) 459–462
Michaelis functions, factorization of, for the binding of a ligand to a molecule with two binding sites (Dixon, H. B. F.) 271–274
Micrococcus flavus N.C.I.B. 8166, structure and properties of lipomannans associated with membranes of, and other micrococci (Powell, D. A., Duckworth, M. & Baddiley, J.) 387–397
Micrococcus lysodeikticus A.T.C.C. 4698 and N.C.I.B. 9278, structure and properties of lipomannans associated with membranes of, and other micrococci (Powell, D. A., Duckworth, M. & Baddiley, J.) 387–397
Micrococcus sodonensis N.C.I.B. 8854, structure and properties of lipomannans associated with membranes of, and other micrococci (Powell, D. A., Duckworth, M. & Baddiley, J.) 387–397
Microsomal fraction, kidney, rabbit, differential effects of temperature on the activity of membrane sodium-plus-potassium ion-dependent adenosine triphosphatase and on the associated phosphatase activity of (Walker, J. A. & Wheeler, K. P.) 439–442
Microsomal fraction, liver, rat, rapid nicotinamide–adenine dinucleotide-linked method for the assay of the activity of uridine diphosphate glucuronyltransferase from, and the substrate specificity of the enzyme (Mulder, G. J. & Van Doorn, A. B. D.) 131–140
Microsomal fraction, salt-gland, dogfish, reversible delipidation of solubilized sodium-plus-potassium ion-dependent adenosine triphosphatase from (Ottolenghi, P.) 61–66
Mitochondria, heart, pig, reversible modification by pyridoxal 5′-phosphate of malate dehydrogenase from (Chen, S.-S. & Engel, P. C.) 297–303
Mitochondria, liver, rat, reaction of choline dehydrogenase from, with some electron acceptors (Barrett, M. C. & Dawson, A. P.) 677–683
Mitochondria, liver, sheep, effects of disulfiram on the activities of aldehyde dehydrogenases from, and cytoplasm (Kitson, T. M.) 407–412
Monkey, capuchin, transport of 25-hydroxycholecalciferol in the plasma of (Hay, A. W. M.) 193–196
Mucosa, small-intestine, calf, affinity-chromatographic purification of alkaline phosphatase from (Brenna, O., Perrella, M., Pace, M. & Pietta, P. G.) 291–296
Muscle, flight, water-bug and dung-beetle, purification of myosin, tropomyosin and α-actinin from (Hammond, K. S. & Goll, D. E.) 189–192
Muscle, heart, rabbit, phosphorylation of light-chain components of myosin from, and red skeletal muscle (Frearson, N. & Perry, S. V.) 99–107
Muscle, skeletal, dogfish, reversible reaction of pyridoxal 5′-phosphate with an essential lysine residue of lactate dehydrogenase M_4 isoenzyme from (Chen, S.-S. & Engel, P. C.) 447–449
Muscle, skeletal, mouse, evaluation by affinity chromatography of the equilibrium constants for the interaction of reduced nicotinamide–adenine dinucleotide with lactate dehydrogenase isoenzymes from, and other tissues (Brinkworth, R. I., Masters, C. J. & Winzor, D. J.) 631–636
Muscle, skeletal, rabbit, effect of pH on the rate of formation and on the equilibrium concentration of the carbanion intermediate formed in the reaction catalysed by fructose 1,6-diphosphate aldolase from (Grazi, E.) 167–172
Muscle, skeletal, rabbit, investigation of substrate-induced changes in subunit interactions in glyceraldehyde 3-phosphate dehydrogenases from baker's yeast and, by measurement of the kinetics and thermodynamics of subunit exchange (Osborne, H. H. & Hollaway, M. R.) 37–45
Muscle, skeletal, rabbit, primary structure of troponin T from, and its interaction with tropomyosin (Jackson, P., Amphlett, G. W. & Perry, S. V.) 85–97
Muscle, skeletal, rabbit, quenched-flow study of the reaction catalysed by creatine kinase from (Engelborghs, Y., Marsh, A. & Gutfreund, H.) 47–50
Muscle, skeletal, red, rabbit, phosphorylation of light-chain components of myosin from, and heart muscle (Frearson, N. & Perry, S. V.) 99–107
Myelin, forebrain, rat, ontogenic study of a myelin derived membrane fraction with higher activity of 2′:3′cyclic nucleotide 3′-phosphohydrolase than that of (Waehneldt, T. V.) 435–437

Myosin, heart-muscle and red-skeletal-muscle, rabbit, phosphorylation of light-chain components of (Frearson, N. & Perry, S. V.) 99–107
Myosin, purification of α-actinin, tropomyosin and, from water-bug and dung-beetle flight muscle (Hammond, K. S. & Goll, D. E.) 189–192

Nasal septum, ox, elecron-microscopic studies on proteoglycans isolated from hyaline cartilage from, and other sources (Thyberg, J., Lohmander, S. & Heinegård, D.) 157–166
Nasal septum, ox, identification of polydispersity of the chondroitin sulphate component in three proteoglycans from cartilage of (Hopwood, J. J. & Robinson, H. C.) 581–594
Nicotiana tabacum, *see* Tobacco
Nicotinamide–adenine dinucleotide, oxidized and reduced, investigation of the changes induced by, in subunit interactions in rabbit skeletal-muscle and baker's-yeast glyceraldehyde 3-phosphate dehydrogenases by measurement of the kinetics and thermodynamics of subunit exchange (Osborne, H. H. & Hollaway, M. R.) 37–45
Nicotinamide–adenine dinucleotide, oxidized, binding of, to *Bacillus stearothermophilus* glyceraldehyde 3-phosphate dehydrogenase (Allen, G. & Harris, J. I.) 747–749
Nicotinamide–adenine dinucleotide, reduced, evaluation by affinity chromatography of the equilibrium constants for the interaction of lactate dehydrogenase isoenzymes from mouse skeletal muscle, heart, liver and kidney with (Brinkworth, R. I., Masters, C. J. & Winzor, D. J.) 631–636
7-Nitrobenzo-2-oxa-1,3-diazole moiety, absolute and selective specificity of the reactions of fluorescent compounds containing, with thiol groups of proteins (Stuchbury, T., Shipton, M., Norris, R., Malthouse, J. P. G., Brocklehurst, K., Herbert, J. A. L., & Suschitzky, H.) 417–432
Nuclei, liver, rat, effects of polyamines and methylglyoxal bis(guanylhydrazone) on the structure of, and deoxyribonucleic acid template activity (Brown, K. B., Nelson, N. F. & Brown, D. G.) 505–512
Nuclei, oviduct, chick, resistance to nuclease of chromatin deoxyribonucleic acid of (Krall, J. F., Socher, S. H., Van, N. T. & O'Malley, B. W.) 497–503
Nucleoside diphosphokinase, activity of, associated with one of the two high-molecular-weight deoxyribonucleic acid polymerases of *Euglena gracilis* (McLennan, A. G. & Keir, H. M.) 239–247
Nucleotides, evidence from multiple-inhibition studies for isosteric and allosteric inhibition by, of the activity of citrate synthases from pig heart, *Bacillus megaterium* and *Pseudomonas aeruginosa* (Harford, S. & Weitzman, P. D. J.) 455–458
Nucleus pulposus, pig, structure of keratan sulphates from, and other sources (Choi, H. U. & Meyer, K.) 543–553

Oestradiol-17β-binding proteins, three, comparison of the specificities of (Laurent, C., de Lauzon, S., Cittanova, N., Nunez, E. & Jayle, M.-F.) 513–518
Oestradiol-17β 3-glucuronide, radioimmunoassay of, and other oestrogen glucuronides (Samarajeewa, P. & Kellie, A. E.) 369–376
Oestradiol-17β, purification and properties of peroxidase induced by, from rat uterus (McNabb, T. & Jellinck, P. H.) 275–279
Oestriol 3-glucuronide, radioimmunoassay of, and other oestrogen glucuronides (Samarajeewa, P. & Kellie, A. E.) 369–376
Oestrogen glucuronides, radioimmunoassay of (Samarajeewa, P. & Kellie, A. E.) 369–376
Oestrogens, purification and properties of peroxidase induced by, from rat uterus (McNabb, T. & Jellinck, P. H.) 275–279
Oestrone 3-glucuronide, radioimmunoassay of, and other oestrogen glucuronides (Samarajeewa, P. & Kellie, A. E.) 369–376
Oligodeoxyribonucleotides, methylated, role of apurinic sites in the resistance of, to degradation by pig spleen exodeoxyribonuclease (Margison, G. P., O'Connor, P. J. & Cornish-Bowden, A.) 249–256
Oplophorus gracilorostris, *see* Shrimp, bioluminescent
Oviduct, chick, resistance to nuclease of chromatin deoxyribonucleic acid from (Krall, J. F., Socher, S. H., Van, N. T. & O'Malley, B. W.) 497–503
Oxyluciferin, purification and properties of, from the bioluminescent shrimp *Oplophorus gracilorostris* and the chemical synthesis of a model luciferin compound biologically active with luciferase from the same organism (Yamaguchi, I.) 9–15

Pancreas, rat, immunochemical relationship between α-amylases of, and liver, serum and parotid gland (Messer, M. & Dean, R. T.) 17–22
Papain, absolute and selective specificity of the reactions of the fluorescent compounds 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole and 4-(*N*-2-aminoethyl 2′-pyridyl disulphide)-7-nitrobenzo-2-oxa-1,3-diazole with thiol groups of, and other proteins (Stuchbury, T., Shipton, M., Norris, R., Malthouse, J. P. G., Brocklehurst, K., Herbert, J. A. L. & Suschitzky, H.) 417–432
Parotid gland, rat, immunochemical relationship between α-amylases of, and liver, serum and pancreas (Messer, M. & Dean, R. T.) 17–22
Penicillin, behaviour of tritium-labelled isopenicillin N and 6-aminopenicillanic acid as precursors of, in an extract of *Penicillium chrysogenum* (Fawcett, P. A., Usher, J. J. & Abraham, E. P.) 741–746
Penicillin N, synthesis of tritium-labelled 6-aminopenicillanic acid, isopenicillin N and (Usher, J. J., Loder, B. & Abraham, E. P.) 729–739
Penicillinase, *Staphylococcus aureus* N.C.I.B. 11195, amino acid sequence of (Ambler, R. P.) 197–218
Penicillium chrysogenum, behaviour of tritium-labelled isopenicillin N and 6-aminopenicillanic acid as potential penicillin precursors in an extract from (Fawcett, P. A., Usher, J. J. & Abraham, E. P.) 741–746
Pepsin, chicken, crystallization of, and of the pepsinogen from which it is derived (Green, M. L.) 763–764
Pepsin, chicken, effects of inhibitors of acid proteinases on the activity of (Llewellin, J. M. & Green, M. L.) 319–326
Pepsinogen, chicken, crystallization of, and of the derived pepsin (Green, M. L.) 763–764
Peptide hormones, nomenclature of (IUPAC–IUB Commission on Biochemical Nomenclature) 1–4

Peroxidase, activities and other properties of, and other enzymes in the cell wall of mesophyll cells of tobacco leaves (Yung, K.-H. & Northcote, D. H.) 141–144
Peroxidase, oestrogen-induced, purification and properties of, from rat uterus (McNabb, T. & Jellinck, P. H.) 275–279
Pertrimethylsilyl methyl glycosides, characterization by gas–liquid chromatography–mass spectrometry and proton-magnetic-resonance spectroscopy of, obtained in the methanolysis of glycoproteins and glycopeptides (Kamerling, J. P., Gerwig, G. J., Vliegenthart, J. F. G. & Clamp, J. R.) 491–495
pH, effect of, on the rate of formation and on the equilibrium concentration of the carbanion intermediate formed in the reaction catalysed by rabbit skeletal-muscle fructose 1,6-diphosphate aldolase (Grazi, E.) 167–172
Phenol, substrate specificity of rat liver microsomal uridine diphosphate glucuronyltransferase towards derivatives of (Mulder, G. J. & Van Doorn, A. B. D.) 131–140
Phenolase, spinach-beet, action of hydrogen peroxide on the hydroxlyation of *p*-coumarate by (Vaughan, P. F. T. & McIntyre, R. J.) 759–762
Phenylalanine, mechanism of the disaggregation by, of polyribosomes in neonatal mouse brain tissue (Taub, F. & Johnson, T. C.) 173–179
Phosphatase, acid, activities and other properties of, and other enzymes in the cell wall of mesophyll cells of tobacco leaves (Yung, K.-H. & Northcote, D. H.) 141–144
Phosphatase, alkaline, affinity-chromatographic purification of, from calf small-intestine mucosa (Brenna, O., Perrella, M., Pace, M. & Pietta, P. G.) 291–296
Phosphatase, differential effects of temperature on the activity of rabbit kidney microsomal membrane sodium-plus-potassium ion-dependent adenosine triphosphatase and on the associated activity of (Walker, J. A. & Wheeler, K. P.) 439–442
Phosphatidylcholine, content of, in pulmonary surfactants from ox, rabbit, rat and sheep (Harwood, J. L., Desai, R., Hext, P., Tetley, T. & Richards, R.) 707–714
Phosphatidylcholine, evidence that the biosynthesis of phosphatidylethanolamine and, is catalysed by a single enzyme in the endoplasmic reticulum of castor-bean endosperm (Lord, J. M.) 451–453
Phosphatidylcholine, physical properties of the phosphonium analogue of, and its suitability as ^{31}P nuclear-magnetic-resonance probe for model and biological membranes (Sim, E., Cullis, P. R. & Richards, R. E.) 555–560
Phosphatidylethanolamine, evidence that the biosynthesis of phosphatidylcholine and, is catalysed by a single enzyme in the endoplasmic reticulum of castor-bean endosperm (Lord, J. M.) 451–453
Phosphofructokinase, purification and properties of, from pig spleen (Hickman, P. E. & Weidemann, M. J.) 327–336
6-Phosphogluconate dehydrogenase, purification and properties of, from rabbit mammary gland (Betts, S. A. & Mayer, R. J.) 263–270
Phosphonium phosphatidylcholine, physical properties of, and its suitability as a ^{31}P nuclear-magnetic-resonance probe for model and biological membranes (Sim, E., Cullis, P. R. & Richards, R. E.) 555–560
Phosvitin, egg-yolk, chicken, oxidation of ferrous iron and the formation of transferrin from apotransferrin catalysed by (Osaki, S., Sexton, R. C., Pascual, E. & Frieden, E.) 519–525
Plasma albumin, *see* Albumin, plasma
Plasma, capuchin-monkey, transport of 25-hydroxycholecalciferol in (Hay, A. W. M.) 193–196
Plasma, human, intramolecular electron transport in ferroxidase from (De Ley, M. & Osaki, S.) 561–566
Plasma membrane, *see* Membrane, plasma
Plasma, ox, further evidence for the lipoprotein nature of arylesterase from (Don, M. M., Masters, C. J. & Winzor, D. J.) 625–630
Polyadenylate, micro method for the assay of ribonucleic acid containing, by gel electrophoresis (Lara, F. J. S. & Okretic, M. C.) 575–580
Polyamines, effects of methylglyoxal bis(guanylhydrazone) and, on rat liver nuclear structure and deoxyribonucleic acid template activity (Brown, K. B., Nelson, N. F. & Brown, D. G.) 505–512
Polyamines, effects of, on the activity of purified baker's-yeast *S*-adenosylmethionine decarboxylase (Pösö, H., Sinervirta, R. & Jänne, J.) 67–73
Polymorphonuclear leucocytes, *see* Leucocytes, polymorphonuclear
Polypeptide chains, characterization of, of ox epidermal α-keratin (Steinert, P. M. & Idler, W. W.) 603–614
Polyribosomes, brain, mouse, neonatal, mechanism of the disaggregation by phenylalanine of (Taub, F. & Johnson, T. C.) 173–179
Polysomes, *see* Polyribosomes
Potassium chloride, stimulation by, and other salts of the activity of rat liver β-galactosidase (Baccion, F. M., Zuretti, M. F. & Pernigotti, L.) 567–573
Protease, *see* Proteinase
Protein, Bence-Jones, λ, identification of, in the urine of guinea pigs with L_2C lymphocytic leukaemia (Stevenson, F. K., Mole, L. E., Raymont, C. M. & Stevenson, G. T.) 751–753
Protein, content of lipid and, in pulmonary surfactants from ox, rabbit, rat and sheep (Harwood, J. L., Desai, R., Hext, P., Tetley, T. & Richards, R.) 707–714
Protein kinase, skeletal-muscle, rabbit, specific, phosphorylation by, of light-chain components of myosin from rabbit heart muscle and red skeletal muscle (Frearson, N. & Perry, S. V.) 99–107
Protein kinases, adenosine 3′:5′-cyclic monophosphate-dependent, properties of, associated with the plasma membrane of ox corpus luteum (Azhar, S. & Menon, K. M. J.) 23–36
Protein, mechanism of the disaggregation by phenylalanine of polyribosomes in neonatal mouse brain tissue and the consequent disruption of the biosynthesis of (Taub, F. & Johnson, T. C.) 173–179
Proteinase, polymorphonuclear-leucocyte, rabbit, evidence for immunological non-identity of rabbit synovial-fibroblast collagenase with (Werb, Z. & Reynolds, J. J.) 665–669
Proteinase (thermomycolin), serine, extracellular, *Malbranchea pulchella* var. *sulfurea*, substrate specificity of (Stevenson, K. J. & Gaucher, G. M.) 527–542
Proteinases, acid, effects of inhibitors of, on the activity of chicken pepsin (Llewellin, J. M. & Green, M. L.) 319–326
Protein–polysaccharides, *see* Proteoglycans

Proteins, absolute and selective specificity of the reactions of the fluorescent compounds 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole and 4-(*N*-2-aminoethyl 2′-pyridyl disulphide)-7-nitrobenzo-2-oxa-1,3-diazole with thiol groups of (Stuchbury, T., Shipton, M., Norris, R., Malthouse, J. P. G., Brocklehurst, K., Herbert, J. A. L. & Suschitzky, H.) 417–432
Proteins, globular, theoretical approach to the binding of amphipathic molecules to (Jones, M. N.) 109–114
Proteins, oestradiol-17β-binding, three, comparison of the specificities of (Laurent, C., de Lauzon, S., Cittanova, N., Nunez, E. & Jayle, M.-F.) 513–518
Proteins, plasma-membrane, corpus luteum, ox, specific, phosphorylation of, by adenosine 3′:5′-cyclic monophosphate-dependent protein kinases associated with the plasma membrane (Azhar, S. & Menon, K. M. J.) 23–36
Proteoglycans, cartilage, nasal-septum, ox, three, identification of polydispersity in the chondroitin sulphate component in (Hopwood, J. J. & Robinson, H. C.) 581–594
Proteoglycans, hyaline-cartilage, guinea-pig and ox, isolated, electron-microscopic studies on (Thyberg, J., Lohmander, S. & Heinegård, D.) 157–166
Pseudomonas aeruginosa, demonstration of two isoaccepting species of lysine transfer ribonucleic acid in Thimmappaya, B. & Cherayil, J. D.) 377–386
Pseudomonas aeruginosa, evidence from multiple-inhibition studies for isosteric and allosteric inhibition by nucleotides of the activity of citrate synthases from, and other sources (Harford, S. & Weitzman, P. D. J.) 455–458
Pseudomonas aeruginosa N.C.T.C. 6750, reaction of carbon monoxide with cytochrome *c*-551 oxidase from (Parr, S. R., Wilson, M. T. & Greenwood, C.) 51–59
Pseudomonas aeruginosa N.C.T.C. 6750, temperature-jump study of the reaction between azurin and cytochrome *c*-551 oxidase from (Brunori, M., Parr, S. R., Greenwood, C. & Wilson, M. T.) 185–188
Pulmonary surfactants, characterization of, from ox, rabbit, rat and sheep (Harwood, J. L., Desai, R., Hext, P., Tetley, T. & Richards, R.) 707–714
Pyridoxal 5′-phosphate, reversible interaction of, with an essential lysine residue of pig heart mitochondrial malate dehydrogenase (Chen, S.-S. & Engel, P. C.) 297–303
Pyridoxal 5′-phosphate, reversible reaction of, with an essential lysine residue of dogfish skeletal-muscle lactate dehydrogenase M_4 isoenzyme (Chen, S.-S. & Engel, P. C.) 447–449
Pyruvate, binding of, to pig heart lactate dehydrogenase and the interconversion of pyruvate-containing ternary complexes (Boland, M. J. & Gutfreund, H.) 715–727

Reticulum, endoplasmic, castor-bean-endosperm, evidence that the biosynthesis of phosphatidylcholine and phosphatidylethanolamine is catalysed by a single enzyme in (Lord, J. M.) 451–453
Rhynchosciara americana, *see* Fly
Rib, guinea-pig, electron-microscopic studies on proteoglycans isolated from hyaline cartilate from, and other sources (Thyberg, J., Lohmander, S. & Heinegård, D.) 157–166
Ribonuclease A, theoretical approach to the binding of amphipathic molecules to, and other globular proteins (Jones, M. N.) 109–114
Ribonucleic acid, messenger, salivary-gland, fly (*Rhynchoscaria americana*)-larva, gel-electrophoretic analysis of (Lara, F. J. S. & Okretic, M. C.) 575–580
Ribonucleic acid, polyadenylate-containing, micro method for the assay of, by gel electrophoresis (Lara, F. J. S. & Okretic, M. C.) 575–580
Ribonucleic acid, transfer, lysine, demonstration of two isoaccepting species of, in *Pseudomonas aeruginosa* (Thimmappaya, B. & Cherayil, J. D.)377–386
Ribosomes, 30S, accumulation of a 30S precursor of, in a mutant of *Escherichia coli* (Markey, F. & Wild, D. G.) 463–465
Ricinus communis, *see* Bean, castor

Saccharomyces cerevisiae, purification and properties of *S*-adenosylmethionine decarboxylase from (Pösö, H., Sinervirta, R. & Jänne, J.) 67–73
Salivary gland, fly (*Rhynchosciara americana*)-larva, gel-electrophoretic analysis of messenger ribonucleic acid from (Lara, F. J. S. & Okretic, M. C.) 575–580
Salivary gland, parotid, rat, immunochemical relation ship between α-amylases of, and liver, serum and pancreas (Messer, M. & Dean, R. T.) 17–22
Salt gland, rectal, dogfish, reversible delipidation of solubilized microsomal sodium-plus-potassium ion-dependent adenosine triphosphatase from (Ottolenghi, P.) 61–66
Serum albumin, *see* Albumin, serum
Serum, interaction of the subtilopeptidase A isoenzyme system with components of, and its importance for quantitative immunoelectrophoresis (Verbruggen, R.) 149–155
Serum, rat, immunochemical relationship between α-amylases of, and liver, pancreas and parotid gland (Messer, M. & Dean, R. T.) 17–22
Shrimp, bioluminescent (*Oplophorus gracilorostris*), purification and properties of oxyluciferin from, and the chemical synthesis of a model luciferin compound biologically active with luciferase from the same organism (Yamaguchi, I.) 9–15
Sialidase, use of affinity chromatography in an improved method for the purification of, from *Clostridium perfringens* (Geisow, M. J.) 181–183
Silver ions, dissociation by, of active *Escherichia coli* β-galactosidase dimer into an inactive monomer (Kaneshiro, C. M., Enns, C. A., Hahn, M. G., Peterson, J. S. & Reithel, F. J.) 433–434
Skeletal muscle, *see* Muscle, skeletal
Skin, ox, characterization of the polypeptide chains of α-keratin from (Steinert, P. M. & Idler, W. W.) 603–614
Small intestine, *see* Intestine, small
Sodium dodecyl sulphate, theoretical approach to the binding of amphipathic molecules such as, to globular proteins (Jones, M. N.) 109–114
Spermine, effects of methylglyoxal bis(guanylhydrazone) and, and other polyamines on rat liver nuclear structure and deoxyribonucleic acid template activity (Brown, K. B., Nelson, N. F. & Brown, D. G.) 505–512
Spleen, pig, purification and properties of phosphofructokinase from (Hickman, P. E. & Weidemann, M. J.) 327–336

Spleen, pig, role of apurinic sites in the resistance of methylated oligodeoxyribonucleotides to degradation by exodeoxyribonuclease from (Margison, G. P., O'Connor, P. J. & Cornish-Bowden, A.) 249–256
Squalus acanthias, *see* Dogfish
Staphylococcus aureus H, influence of alanine ester residues on the binding of magnesium ions to teichoic acids from the cell wall of (Lambert, P. A., Hancock, I. C. & Baddiley, J.) 671–676
Staphylococcus aureus N.C.I.B. 11195, amino acid sequence of penicillinase from (Ambler, R. P.) 197–218
Subtilopeptidase A, interaction of the isoenzyme system of, with serum components and its importance for quantitative immunoelectrophoresis (Verbruggen, R.) 149–155
Supernatant fraction, *see* Cytosol
Surfactants, pulmonary, characterization of, from ox, rabbit, rat and sheep (Harwood, J. L., Desai, R., Hext, P., Tetley, T. & Richards, R.) 707–314
Synovial fibroblasts, rabbit, cultured, immunochemical studies with a specific antiserum to collagenase from (Werb, Z. & Reynolds, J. J.) 655–663
Synovial fibroblasts, rabbit, cultured, immunological identity of the collagenase from, with the enzymes released from other rabbit cells and tissues in normal and pathological conditions (Werb, Z. & Reynolds, J. J.) 665–669
Synovial fibroblasts, rabbit, cultured, purification and properties of a specific collagenase from (Werb, Z. & Reynolds, J. J.) 645–653

Teichoic acid, poly(glucosylglycerol phosphate), characterization of, in the cell wall of *Bacillus stearothermophilus* B 65 (Anderson, A. J. & Archibald, A. R.) 115–120
Teichoic acids, influence of alanine ester residues on the binding of magnesium ions to, from *Staphylococcus aureus* H cell wall (Lambert, P. A., Hancock, I. C. & Baddiley, J.) 671–676
Temperature, differential effects of, on the activity of rabbit kidney microsomal membrane sodium-plus-potassium ion-dependent adenosine triphosphatase and on the associated phosphatase activity (Walker, J. A. & Wheeler, K. P.) 439–442
Tetraethylthiuram disulphide (disulfiram), effects of, on the activities of sheep liver cytoplasmic and mitochondrial aldehyde dehydrogenases (Kitson, T. M.) 407–412
Thermomycolase, *see* Thermomycolin
Thermomycolin, *Malbranchea pulchella* var. *sulfurea*, extracellular, substrate specificity of (Stevenson, K. J. & Gaucher, G. M.) 527–542
Thiol groups, protein, absolute and selective specificity of the reactions of the fluorescent compounds 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole and 4-(*N*-2-aminoethyl 2′-pyridyl disulphide)-7-nitrobenzo-2-oxa-1,3-diazole with (Stuchbury, T., Shipton, M., Norris, R., Malthouse, J. P. G., Brocklehurst, K., Herbert, J. A. L. & Suschitzky, H.) 417–432
Thy-1 glycoprotein, purification and characterization of, from rat brain membranes (Barclay, A. N., Letarte-Muirhead, M. & Williams, A. F.) 699–706
Thy-1 glycoprotein, purification and characterization of, from rat thymocyte membranes (Letarte-Muirhead, M., Barclay, A. N. & Williams, A. F.) 685–697
Thymocytes, rat, purification and characterization of Thy-1 glycoprotein from membranes of (Letarte-Muirhead, M., Barclay, A. N. & Williams, A. F.) 685–697
Thymus gland, rat, purification and characterization of Thy-1 glycoprotein from cell membranes of (Letarte-Muirhead, M., Barclay, A. N. & Williams, A. F.) 685–697
Tobacco (*Nicotiana tabacum*) leaves, activities and other properties of enzymes in the cell wall of mesophyll cells of (Yung, K.-H. & Northcote, D. H.) 141–144
Trachea, ox, electron-microscopic studies on proteoglycans isolated from hyaline cartilage from, and other sources (Thyberg, J., Lohmander, S. & Heinegård, D.) 157–166
Transfer ribonucleic acid, *see* Ribonucleic acid, transfer
Transferrin, apo-, oxidation of ferrous iron and the formation of transferrin from, catalysed by chicken egg-yolk phosvitin (Osaki, S., Sexton, R. C., Pascual, E. & Freiden, E.) 519–525
Transferrin, oxidation of ferrous iron and the formation of, from apotransferrin catalysed by chicken egg-yolk phosvitin (Osaki, S., Sexton, R. C., Pascual, E. & Freiden, E.) 519–525
Trichomonas foetus, purification and properties of β-*N*-acetylhexosaminidase from (Edwards, R. G., Thomas, P. & Westwood, J. H.) 145–148
Triticum vulgare, *see* Wheat
Tritium isotope (^{3}H), semi-synthesis of pig insulin labelled with, with a specific radioactivity of up to 20 curies per millimole (Halban, P. A. & Offord, R. E.) 219–225
Tritium isotope (^{3}H), synthesis of isopenicillin N, penicillin N and 6-aminopenicillanic acid labelled with (Usher, J. J., Loder, B. & Abraham, E. P.) 729–739
Tropomyosin, purification of α-actinin, myosin and, from water-bug and dung-beetle flight muscle (Hammond, K. S. & Goll, D. E.) 189–192
Tropomyosin, skeletal-muscle, rabbit, primary structure of troponin T and its interaction with (Jackson, P., Amphlett, G. W. & Perry, S. V.) 85–97
Troponin C, skeletal-muscle, rabbit, primary structure of troponin T and its interaction with, and tropomyosin (Jackson, P., Amphlett, G. W. & Perry, S. V.) 85–97
Troponin T, skeletal-muscle, rabbit, primary structure of, and its interaction with tropomyosin (Jackson, P., Amphlett, G. W. & Perry, S. V.) 85–97
Tyrosinase, spinach-beet, action of hydrogen peroxide on the hydroxylation of *p*-coumarate by (Vaughan, P. F. T. & McIntyre, R. J.) 759–762
Tyrosine, transamination of, by rat liver, brain and small-intestine kynurenine–2-oxoglutarate aminotransferase (Noguchi, T., Minatogawa, Y., Okuno, E., Nakatani, M., Morimoto, M. & Kido, R.) 399–406

Ubiquinone-2, reaction of rat liver mitochondrial choline dehydrogenase with, and other electron acceptors (Barrett, M. C. & Dawson, A. P.) 677–683
Uridine diphosphate glucuronyltransferase, microsomal, liver, rat, rapid nicotinamide–adenine dinucleotide-linked method for the assay of the activity of, and the substrate specificity of the enzyme (Mulder, G. J. & Van Doorn, A. B. D.) 131–140

Urine, identification of a λ Bence-Jones protein in, of guinea pigs with L_2C lymphocytic leukaemia (Stevenson, F. K., Mole, L. E., Raymont, C. M. & Stevenson, G. T.) 751–753

Urine, pregnancy, radioimmunoassay of oestriol 3-glucuronide and other oestrogen glucuronides in (Samarajeewa, P. & Kellie, A. E.) 369–376

Uterus, rat, purification and properties of oestrogen-induced peroxidase from (McNabb, T. & Jellinck, P. H.) 275–279

Vitamin D_3, *see* Cholecalciferol

Wall, cell, *see* Cell wall

Water bug (*Lethocerus cordofanus*), purification of myosin, tropomyosin and α-actinin from the flight muscle of (Hammond, K. S. & Goll, D. E.) 189–192

Wheat (*Triticum vulgare*) germ, interaction of *N*-(3-carboxypropionyl)aminodecyl-Sepharose with aspartate carbamoyltransferase from (Yon, R. J. & Simmonds, R. J.) 281–290

Yeast, baker's, investigation of substrate-induced changes in subunit interactions in glyceraldehyde 3-phosphate dehydrogenases from rabbit skeletal muscle and, by measurement of the kinetics and thermodynamics of subunit exchange (Osborne, H. H. & Hollaway, M. R.) 37–45

Yeast, baker's (*Saccharomyces cerevisiae*), purification and properties of *S*-adenosylmethionine decarboxylase from (Pösö, H., Sinervirta, R. & Jänne, J.) 67–73